



POC5 Rabbit mAb

Catalog No	YP-rAb-17682
Isotype	IgG
Reactivity	Human
Applications	WB,IHC
Gene Name	POC5;C5orf37
Protein Name	Centrosomal protein POC5;Protein of centriole 5;hPOC5;
Purification Process	Protein A
Specificity	Endogenous
Formulation	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source	Monoclonal, Rabbit,IgG
Dilution	WB 1:1000-5000; IHC 1:100-300 Note: For IHC, we suggest antigen retrieval with TE buffer pH 9.0
Concentration	0.5 mg/ml
Purity	≥90%
Storage Stability	-15° C to -25° C/1 year(Do not lower than -25° C)
Synonyms	POC5 ; C5orf37 ; Centrosomal protein POC5 ; Protein of centriole 5 ; hPOC5 ;
Observed Band	63kD
Calculated Molecular Weight	
Cell Pathway	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome.centriole. Note=Localized to the distal portion of centrioles.
Tissue Specificity	
Function	Essential for the assembly of the distal half of centrioles, required for centriole elongation. {ECO:0000269 PubMed:19349582}.
Background	Predicted to be involved in eye photoreceptor cell development. Located in centrosome; cytosol; and nucleoplasm. [provided by Alliance of Genome Resources, Feb 2025]
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





HEK-293

kDa

200-

140-

100-

70-

55-

40-

35-

25-

20-

15-

10-



POC5

Western Blot analysis of HEK-293 whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-POC5 rabbit mAb. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody.





杭州臻优品生物科技有限公司

热销产品:

蛋白、一抗、抗体对、ELISA试剂盒、生化试剂盒
CCK8试剂盒、QPCR检测试剂盒

检测服务:

ELISA检测及定制服务 | 生化检测 | PCR、QPCR检测 | WB检测
ICO-IP检测 | 切片 | 染色 | 免疫组化 | 免疫荧光 | 透射电镜全套
| 宏基因组、转录组、基因组、蛋白组、代谢组测序



关注官网



关注客服