



HBO1 Rabbit mAb

Catalog No	YP-rAb-17774
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB,IHC,IF,IP,ELISA
Gene Name	KAT7
Protein Name	Histone acetyltransferase KAT7
Purification Process	Protein A
Specificity	Endogenous
Formulation	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source	Monoclonal, Rabbit,IgG
Dilution	IHC 1:200-1:1000; WB 1:2000-1:10000; IF 1:200-1:1000; ELISA 1:5000-1:20000; IP 1:50-1:200; Note: For IHC, we suggest antigen retrieval with TE buffer pH 9.0
Concentration	0.5 mg/ml
Purity	≥90%
Storage Stability	-15° C to -25° C/1 year(Do not lower than -25° C)
Synonyms	KAT7 ; HBO1 ; HBOa ; MYST2 ; Histone acetyltransferase KAT7 ; Histone acetyltransferase binding to ORC1 ; Lysine acetyltransferase 7 ; MOZ ; YBF2/SAS3, SAS2 and TIP60 protein 2 ; MYST-2
Observed Band	71kD
Calculated Molecular Weight	71kD
Cell Pathway	Nucleus . Chromosome . Chromosome, centromere . Cytoplasm, cytosol . Associates with replication origins specifically during the G1 phase of the cell cycle (PubMed:18832067, PubMed:20129055). Localizes to transcription start sites (PubMed:21753189, PubMed:24065767). Localizes to ultraviolet-induced DNA damage sites following phosphorylation by ATR (PubMed:28719581). Localizes to centromeres in G1 phase (PubMed:27270040). .
Tissue Specificity	Ubiquitously expressed, with highest levels in testis.
Function	Catalytic activity:Acetyl-CoA + histone = CoA + acetylhistone.,Domain:The C2HC-type zinc finger is required for interaction with MCM2 and ORC1L.,Domain:The N-terminus is involved in transcriptional repression, while the C-terminus mediates AR-interaction.,Function:Component of the HBO1 complex which has a histone H4-specific acetyltransferase activity, a reduced activity toward histone H3 and is responsible for the bulk of histone H4 acetylation in vivo. Through chromatin acetylation it may regulate DNA replication and act as

杭州臻优品生物科技有限公司

热销产品:

蛋白、一抗、抗体对、ELISA试剂盒、生化试剂盒
CCK8试剂盒、QPCR检测试剂盒

检测服务:

ELISA检测及定制服务 | 生化检测 | PCR、QPCR检测 | WB检测
ICO-IP检测 | 切片 | 染色 | 免疫组化 | 免疫荧光 | 透射电镜全套
| 宏基因组、转录组、基因组、蛋白组、代谢组测序



关注官网



关注客服



a coactivator of TP53-dependent transcription. Specifically represses AR-mediated transcription.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the MYST (SAS/MOZ) family.,similarity:Contains 1 C2HC-type zinc finger.,subunit:Component of the HBO1 complex composed at least of ING4 or ING5, MYTS2/HBO1, EAF6, and one of PHF15, PHF16 and PHF17. Interacts with MCM2 and ORC1L. Interacts with the androgen receptor (AR) in the presence of dihydrotestosterone.,tissue specificity:Ubiquitously expressed, with highest levels in testis.,

Background


catalytic activity:Acetyl-CoA + histone = CoA + acetylhistone.,domain:The C2HC-type zinc finger is required for interaction with MCM2 and ORC1L.,domain:The N-terminus is involved in transcriptional repression, while the C-terminus mediates AR-interaction.,function:Component of the HBO1 complex which has a histone H4-specific acetyltransferase activity, a reduced activity toward histone H3 and is responsible for the bulk of histone H4 acetylation in vivo. Through chromatin acetylation it may regulate DNA replication and act as a coactivator of TP53-dependent transcription. Specifically represses AR-mediated transcription.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the MYST (SAS/MOZ) family.,similarity:Contains 1 C2HC-type zinc finger.,subunit:Component of the HBO1 complex composed at least of ING4 or ING5, MYTS2/HBO1, EAF6, and one of PHF15, PHF16 and PHF17. Interacts with MCM2 and ORC1L. Interacts with the androgen receptor (AR) in the presence of dihydrotestosterone.,tissue specificity:Ubiquitously expressed, with highest levels in testis.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-HBO1 antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: C6 Lane 2: NIH-3T3 Lane 3: HeLa Lane 4: Jurkat Predicted band size: 71kDa Observed band size: 71kDa

