



# Fibrillarin Rabbit mAb

|                                    |  |
|------------------------------------|--|
| <b>Catalog No</b>                  | YP-rAb-18064   |
| <b>Isotype</b>                     | IgG  |
| <b>Reactivity</b>                  | Human,Mouse,Rat  |
| <b>Applications</b>                | WB,IHC,IF,ELISA  |
| <b>Gene Name</b>                   | FBL FIB1 FLRN  |
| <b>Protein Name</b>                | rRNA 2'-O-methyltransferase fibrillarin (34 kDa nucleolar scleroderma antigen)   |
| <b>Purification Process</b>        | Protein A  |
| <b>Specificity</b>                 | Endogenous   |
| <b>Formulation</b>                 | PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA   |
| <b>Source</b>                      | Monoclonal, Rabbit,IgG   |
| <b>Dilution</b>                    | IHC 1:200-1000; WB 1:1000-5000; IF 1:200-1000; ELISA 1:5000-20000; Note: For IHC, we suggest antigen retrieval with TE buffer pH 9.0   |
| <b>Concentration</b>               | 0.5 mg/ml  |
| <b>Purity</b>                      | ≥90%   |
| <b>Storage Stability</b>           | -15° C to -25° C/1 year(Do not lower than -25° C)  |
| <b>Synonyms</b>                    |  |
| <b>Observed Band</b>               | 34kD   |
| <b>Calculated Molecular Weight</b> | 34kD   |
| <b>Cell Pathway</b>                | Nucleus  |
| <b>Tissue Specificity</b>          |  |
| <b>Function</b>                    | S-adenosyl-L-methionine-dependent methyltransferase that has the ability to methylate both RNAs and proteins . Involved in pre-rRNA processing by catalyzing the site-specific 2'-hydroxyl methylation of ribose moieties in pre-ribosomal RNA . Site specificity is provided by a guide RNA that base pairs with the substrate (By similarity). Methylation occurs at a characteristic distance from the sequence involved in base pairing with the guide RNA (By similarity). Probably catalyzes 2'-O-methylation of U6 snRNAs in box C/D RNP complexes . U6 snRNA 2'-O-methylation is required for mRNA splicing fidelity . Also acts as a protein methyltransferase by mediating methylation of 'Gln-105' of histone H2A (H2AQ104me), a modification that impairs binding of the FACT complex and is specifically present at 35S ribosomal DNA locus . |





## Background


This gene product is a component of a nucleolar small nuclear ribonucleoprotein (snRNP) particle thought to participate in the first step in processing preribosomal RNA. It is associated with the U3, U8, and U13 small nuclear RNAs and is located in the dense fibrillar component (DFC) of the nucleolus. The encoded protein contains an N-terminal repetitive domain that is rich in glycine and arginine residues, like fibrillarins in other species. Its central region resembles an RNA-binding domain and contains an RNP consensus sequence. Antisera from approximately 8% of humans with the autoimmune disease scleroderma recognize fibrillarins. [provided by RefSeq, Jul 2008]

## matters needing attention


Avoid repeated freezing and thawing!

## Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the primary antibody was used at 4°C, over night with a 1:2500 dilution. The Dylight 800-conjugated Goat anti-Rabbit antibody




Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-Fibrillarins antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: 3T3-L1 Lane 2: A172 Lane 3: C6 Lane 4: K562 Predicted band size: 34kDa Observed band size: 34kDa




Mouse testis was stained with Anti-Fibrillarins rabbit antibody





Rat liver was stained with anti-Fibrillarlin rabbit antibody



Human liver was stained with anti-Fibrillarlin rabbit antibody

