



C9orf167 Rabbit mAb

Catalog No	YP-rAb-18488
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB
Gene Name	TOR4A,C9orf167
Protein Name	Torsin-4A (Torsin family 4 member A)
Purification Process	Protein A
Specificity	Endogenous
Formulation	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source	Monoclonal, Rabbit,IgG
Dilution	WB 1:1000-1:5000;
Concentration	0.5 mg/ml
Purity	≥90%
Storage Stability	-15° C to -25° C/1 year(Do not lower than -25° C)
Synonyms	C9orf167
Observed Band	47kD
Calculated Molecular Weight	47kD
Cell Pathway	SUBCELLULAR LOCATION: Membrane {ECO:0000305}; Single-pass membrane protein {ECO:0000305}.
Tissue Specificity	
Function	
Background	Predicted to enable ATP binding activity and ATP hydrolysis activity. Predicted to be located in extracellular region and platelet alpha granule lumen. Predicted to be active in endoplasmic reticulum lumen and nuclear envelope. [provided by Alliance of Genome Resources, Apr 2025]
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

杭州臻优品生物科技有限公司

热销产品:

蛋白、一抗、抗体对、ELISA试剂盒、生化试剂盒
CCK8试剂盒、QPCR检测试剂盒

检测服务:

ELISA检测及定制服务 | 生化检测 | PCR、QPCR检测 | WB检测
ICO-IP检测 | 切片 | 染色 | 免疫组化 | 免疫荧光 | 透射电镜全套
| 宏基因组、转录组、基因组、蛋白组、代谢组测序



关注官网



关注客服



Western blot analysis of lysates from Hela cell, primary antibody was diluted at 1:1000, 4° over night, Dylight 800 secondary antibody was diluted at 1:10000, 37° 1hour.

