



# UCHL5IP Mouse mAb

<b>Catalog No</b>	YP-mAb-19326
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	HAUS7 UCHL5IP UIP1
<b>Protein Name</b>	HAUS augmin-like complex subunit 7 (26S proteasome-associated UCH37-interacting protein 1) (UCHL5-interacting protein) (X-linked protein STS1769)
<b>Immunogen</b>	Synthesized peptide derived from human UCHL5IP
<b>Specificity</b>	This antibody detects endogenous levels of UCHL5IP at Human, Mouse
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Monoclonal,Mouse,IgG
<b>Purification</b>	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Calculated Molecular Weight</b>	39kD
<b>Cell Pathway</b>	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm, cytoskeleton, spindle . Localizes to interphase centrosomes and to mitotic spindle microtubules. .
<b>Tissue Specificity</b>	Detected in spleen, thymus, testis, ovary, small intestine and colon, with highest levels of expression in testis and ovary.
<b>Function</b>	Contributes to mitotic spindle assembly, maintenance of centrosome integrity and completion of cytokinesis as part of the HAUS augmin-like complex.
<b>Background</b>	
<b>matters needing attention</b>	Avoid repeated freezing and thawing!



### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

### Products Images