



AhR Rabbit mAb

Catalog No	YP-rAb-16924
Isotype	IgG
Reactivity	Human
Applications	WB,IF,ELISA
Gene Name	AHR BHLHE76
Protein Name	Aryl hydrocarbon receptor
Purification Process	Protein A
Specificity	Endogenous
Formulation	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source	Monoclonal, Rabbit,IgG
Dilution	WB 1:2000-1:10000; IF 1:200-1:1000; ELISA 1:5000-1:20000;
Concentration	0.5 mg/ml
Purity	≥90%
Storage Stability	-15 ° C to -25 ° C/1 year(Do not lower than -25 ° C)
Synonyms	AHR ; BHLHE76 ; Aryl hydrocarbon receptor ; Ah receptor ; AhR ; Class E basic helix-loop-helix protein 76 ; bHLHe76 ; AHRR ; BHLHE77 ; KIAA1234 ; Aryl hydrocarbon receptor repressor ; AhR repressor ; AhRR ; Class E basic helix-loop-helix protein 77 ; bHL
Observed Band	110kD
Calculated Molecular Weight	96kD
Cell Pathway	Cytoplasm . Nucleus . Initially cytoplasmic; upon binding with ligand and interaction with a HSP 90, it translocates to the nucleus__
Tissue Specificity	Expressed in all tissues tested including blood, brain, heart, kidney, liver, lung, pancreas and skeletal muscle . Expressed in retinal photoreceptors (PubMed:29726989).
Function	Ligand-activated transcriptional activator. Binds to the XRE promoter region of genes it activates. Activates the expression of multiple phase I and II xenobiotic chemical metabolizing enzyme genes (such as the CYP1A1 gene). Mediates biochemical and toxic effects of halogenated aromatic hydrocarbons. Involved in cell-cycle regulation. Likely to play an important role in the development and maturation of many tissues.,induction:Induced or repressed by TGF-beta and dioxin in a cell-type specific fashion. Repressed by cAMP, retinoic acid, and TPA.,similarity:Contains 1 basic helix-loop-helix (bHLH)





domain.,similarity:Contains 1 PAC (PAS-associated C-terminal)
domain.,similarity:Contains 2 PAS (PER-ARNT-SIM) domains.,subcellular
location:Initially cytoplasmic; upon binding with ligand and interaction with a
HSP90, it translocates to the nucleus.,subunit:Binds MYBBP1A (By similarity) .
Efficient DNA binding requires dimerization with another bHLH protein. In the
nucleus, heterodimer of AHR and ARNT. Interacts with coactivators including
SRC-1, RIP140 and NOCA7, and with the corepressor SMRT. Interacts with
NEDD8 and IVNS1ABP.,tissue specificity:Expressed in all tissues tested including
blood, brain, heart, kidney, liver, lung, pancreas and skeletal muscle.,

Background


The protein encoded by this gene is a ligand-activated helix-loop-helix transcription factor involved in the regulation of biological responses to planar aromatic hydrocarbons. This receptor has been shown to regulate xenobiotic-metabolizing enzymes such as cytochrome P450. Before ligand binding, the encoded protein is sequestered in the cytoplasm; upon ligand binding, this protein moves to the nucleus and stimulates transcription of target genes. [provided by RefSeq, Sep 2015],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-AhR antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: HepG2 Predicted band size: 96kDa Observed band size: 110kDa

