



# E-cadherin Rabbit mAb

|                             |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
|-----------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Catalog No                  | YP-rAb-17108                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| Isotype                     | IgG                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Reactivity                  | Human,Mouse                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Applications                | WB,IHC,IF,ELISA                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Gene Name                   | CDH1                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Protein Name                | Cadherin-1                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Purification Process        | Protein A                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Specificity                 | Endogenous                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                  |
| Formulation                 | PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Source                      | Monoclonal, Rabbit,IgG                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Dilution                    | IHC 1:200-1:1000; WB 1:2000-1:10000; IF 1:200-1:1000; ELISA 1:5000-1:20000;<br>Note: For IHC, we suggest antigen retrieval with TE buffer pH 9.0                                                                                                                                                                                                                                                                                                                                                                                                                            |
| Concentration               | 0.5 mg/ml                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Purity                      | ≥90%                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Storage Stability           | -15 ° C to -25 ° C/1 year(Do not lower than -25 ° C)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Synonyms                    | E-cadherin;Cadherin-1;CDH1 ; CDHE ; UVO ; Cadherin-1 ; CAM 120/80 ; Epithelial cadherin ; E-cadherin ; Uvomorulin ; CD antigen CD324;CDH1                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Observed Band               | 125kD                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Calculated Molecular Weight | 97kD                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Cell Pathway                | Cell junction, adherens junction . Cell membrane ; Single-pass type I membrane protein. Endosome. Golgi apparatus, trans-Golgi network. Colocalizes with DLGAP5 at sites of cell-cell contact in intestinal epithelial cells. Anchored to actin microfilaments through association with alpha-, beta- and gamma-catenin . Sequential proteolysis induced by apoptosis or calcium influx, results in translocation from sites of cell-cell contact to the cytoplasm. Colocalizes with RAB11A endosomes during its transport from the Golgi apparatus to the plasma membrane. |
| Tissue Specificity          | Non-neural epithelial tissues.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |
| Function                    | Disease:Defects in CDH1 are a cause of gastric cancer [MIM:137215]; also known as hereditary familial diffuse gastric cancer (HDGC).,Disease:Defects in CDH1 are a cause of susceptibility to endometrial cancer [MIM:608089].,Disease:Defects in CDH1 are associated with ovarian cancer [MIM:167000]. Ovarian cancer is the leading cause of death from gynecologic malignancy. It is characterized by advanced presentation with loco-regional                                                                                                                           |





dissemination in the peritoneal cavity and the rare incidence of visceral metastases. These typical features relate to the biology of the disease, which is a principal determinant of outcome. Disease: Defects in CDH1 are involved in dysfunction of the cell-cell adhesion system, triggering cancer invasion (gastric, breast, ovary, endometrium and thyroid) and metastasis. Function: Cadherins are calcium dependent cell adhesion proteins. Function: Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. CDH1 is involved in mechanisms regulating cell-cell adhesions, mobility and proliferation of epithelial cells. Has a potent invasive suppressor role. It is a ligand for integrin alpha-E/beta-7. Function: E-Cad/CTF2 promotes non-amyloidogenic degradation of Aβeta precursors. Has a strong inhibitory effect on APP C99 and C83 production. online information: E-cadherin entry, PTM: During apoptosis or with calcium influx, cleaved by a membrane-bound metalloproteinase (ADAM10), PS1/gamma-secretase and caspase-3 to produce fragments of about 38 kDa (E-CAD/CTF1), 33 kDa (E-CAD/CTF2) and 29 kDa (E-CAD/CTF3), respectively. Processing by the metalloproteinase, induced by calcium influx, causes disruption of cell-cell adhesion and the subsequent release of beta-catenin into the cytoplasm. The residual membrane-tethered cleavage product is rapidly degraded via an intracellular proteolytic pathway. Cleavage by caspase-3 releases the cytoplasmic tail resulting in disintegration of the actin microfilament system. The gamma-secretase-mediated cleavage promotes disassembly of adherens junctions. similarity: Contains 5 cadherin domains. subcellular location: Colocalizes with DLGAP5 at sites of cell-cell contact in intestinal epithelial cells. Anchored to actin microfilaments through association with alpha-, beta- and gamma-catenin. Sequential proteolysis induced by apoptosis or calcium influx, results in translocation from sites of cell-cell contact to the cytoplasm. subunit: Homodimer; disulfide-linked. Interacts directly, via the cytoplasmic domain, with CTNNB1 or JUP to form the PSEN1/cadherin/catenin adhesion complex which connects to the actin skeleton through the actin binding of alpha-catenin. Interaction with PSEN1, cleaves CDH1 resulting in the disassociation of cadherin-based adherens junctions (CAJs). Interacts with AJAP1, CTNND1 and DLGAP5. tissue specificity: Non-neural epithelial tissues.

## Background

This gene encodes a classical cadherin of the cadherin superfamily. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate the mature glycoprotein. This calcium-dependent cell-cell adhesion protein is comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. Mutations in this gene are correlated with gastric, breast, colorectal, thyroid and ovarian cancer. Loss of function of this gene is thought to contribute to cancer progression by increasing proliferation, invasion, and/or metastasis. The ectodomain of this protein mediates bacterial adhesion to mammalian cells and the cytoplasmic domain is required for internalization. This gene is present in a gene cluster with other members of the cadherin family on chromosome 16. [provided by RefSeq, Nov 2015]


## matters needing attention

Avoid repeated freezing and thawing!


## Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-E-cadherin antibody. The HRP-conjugated Goat anti-Rabbit IgG (H + L) antibody was used to detect the antibody. Lane 1: A549 Lane 2: 4T1 Predicted band size: 97kDa Observed band size: 125kDa



Human liver was stained with anti-E-cadherin Rabbit antibody

