



PI3 Kinase Class 3 Rabbit mAb

Catalog No	YP-rAb-17534
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB,IHC,IF,ELISA
Gene Name	PIK3C3 VPS34
Protein Name	PI 3 Kinase Class 3
Purification Process	Protein A
Specificity	Endogenous
Formulation	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source	Monoclonal, Rabbit,IgG
Dilution	IHC 1:200-1:1000; WB 1:2000-1:10000; IF 1:200-1:1000; ELISA 1:5000-1:20000; Note: For IHC, we suggest antigen retrieval with TE buffer pH 9.0
Concentration	0.5 mg/ml
Purity	≥90%
Storage Stability	-15 ° C to -25 ° C/1 year(Do not lower than -25 ° C)
Synonyms	Phosphatidylinositol 3-kinase catalytic subunit type 3 ; PI3-kinase type 3 ; PI3K type 3 ; PtdIns-3-kinase type 3 ; Phosphatidylinositol 3-kinase p100 subunit ; Phosphoinositide-3-kinase class 3 ; hVps34 ;
Observed Band	102kD
Calculated Molecular Weight	102kD
Cell Pathway	Midbody . Late endosome . Cytoplasmic vesicle, autophagosome . As component of the p13K complex I localized to pre-autophagosome structures . As component of the PI3 K complex II localized predominantly to endosomes (PubMed: 14617358). Localizes also to discrete punctae along the ciliary axoneme and to the base of the ciliary axoneme (By similarity). .
Tissue Specificity	Ubiquitously expressed, with a highest expression in skeletal muscle.
Function	Catalytic activity:ATP + 1-phosphatidyl-1D-myo-inositol = ADP + 1-phosphatidyl-1D-myo-inositol 3-phosphate.,cofactor:Manganese.,Function:Catalytic subunit of the PI3K complex. Involved in the transport of lysosomal enzyme precursors to lysosomes.,similarity:Belongs to the PI3/PI4-kinase family.,similarity:Contains 1 PI3K/PI4K domain.,subunit:Probably forms a complex with AMBRA1 and BECN1 (By similarity). Heterodimer. This subunit, part of a complex composed of regulatory and catalytic subunits, associates with regulatory subunit





PIK3R4.,tissue specificity:Ubiquitously expressed, with a highest expression in skeletal muscle.,

Background


catalytic activity:ATP + 1-phosphatidyl-1D-myo-inositol = ADP + 1-phosphatidyl-1D-myo-inositol 3-phosphate.,cofactor:Manganese.,function:Catalytic subunit of the PI3K complex. Involved in the transport of lysosomal enzyme precursors to lysosomes.,similarity:Belongs to the PI3/PI4-kinase family.,similarity:Contains 1 PI3K/PI4K domain.,subunit:Probably forms a complex with AMBRA1 and BECN1 (By similarity). Heterodimer. This subunit, part of a complex composed of regulatory and catalytic subunits, associates with regulatory subunit PIK3R4.,tissue specificity:Ubiquitously expressed, with a highest expression in skeletal muscle.,

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the primary antibody was used at 4~C, over night with a 1:5000 dilution . The Dylight 800-conjugated Goat anti-Rabbit antibody




Human liver was stained with anti-PI3 Kinase Class 3 rabbit antibody




Human gastric adenocarcinoma was stained with anti-PI3 Kinase Class 3 rabbit antibody





Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-PI3 Kinase Class 3 antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: Rat brain Lane 2: Mouse brain Lane 3: U-87 MG Predicted band size: 102kDa Observed band size: 102kDa



Immunofluorescence analysis of HEK293. Picture A: PI3 Kinase Class 3 antibody (red). Picture B: DAPI (blue). Picture C: Merge of A+B

