



Egr 1/2 Rabbit pAb

Catalog No	YP-Ab-19481
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB、IHC、IF、ELISA
Gene Name	EGR1/2
Immunogen	The antiserum was produced against synthesized peptide derived from human EGR1/2. AA range:371-420
Specificity	Egr1/2 Polyclonal Antibody detects endogenous levels of Egr1/2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000; IHC 1:100-1:300; IF 1:200-1:1000; ELISA 1:20000; Not yet tested in other applications. Note: For IHC, we suggest antigen retrieval with TE buffer pH 9.0 (Cat#YS0004)
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-15°C to -25°C/1 year(Do not lower than -25°C)
Synonyms	EGR1/2;Early growth response protein 1/2;EGR1 ; KROX24 ; ZNF225 ; Early growth response protein 1 ; EGR-1 ; AT225 ; Nerve growth factor-induced protein A ; NGFI-A ; Transcription factor ETR103 ; Transcription factor Zif268 ; Zinc finger protein 225 ; Zinc finger protein Krox-24 ; EGR2 ; KROX20
Observed Band	50kD
Calculated Molecular Weight	
Cell Pathway	Nucleus . Cytoplasm .
Tissue Specificity	Detected in neutrophils (at protein level) .
Function	Function:Transcriptional regulator. Recognizes and binds to the DNA sequence 5'-CGCCCCCGC-3' (EGR-site) . Activates the transcription of target genes whose products are required for mitogenesis and differentiation. ,induction:By growth factors. ,similarity:Belongs to the EGR C2H2-type zinc-finger protein family. ,similarity:Contains 3 C2H2-type zinc fingers. ,
Background	The protein encoded by this gene belongs to the EGR family of C2H2-type zinc-finger proteins. It is a nuclear protein and functions as a transcriptional regulator. The products of target genes it activates are required for differentiation and mitogenesis. Studies suggest this is a cancer suppressor gene. [provided by



RefSeq , Dec 2014] ,


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HeLa and Jurkat cells, using EGR1/2 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Tris-EDTA, pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200 (4°C overnight). 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).