



EP2 Monoclonal Antibody

Catalog No	YP-mAb-13214
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	PTGER2
Immunogen	The antiserum was produced against synthesized peptide derived from human PE2R2. AA range:261-310
Specificity	EP2 Monoclonal Antibody detects endogenous levels of EP2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	EP2;PTGER2;Prostaglandin E2 receptor EP2 subtype;PGE receptor EP2 subtype;PGE2 receptor EP2 subtype;Prostanoid EP2 receptor
Observed Band	40kD
Calculated Molecular Weight	40kD
Cell Pathway	Cell membrane; Multi-pass membrane protein.
Tissue Specificity	Placenta and lung.
Function	function:Receptor for prostaglandin E2 (PGE2). The activity of this receptor is mediated by G(s) proteins that stimulate adenylate cyclase. The subsequent raise in intracellular cAMP is responsible for the relaxing effect of this receptor on smooth muscle.,similarity:Belongs to the G-protein coupled receptor 1 family., tissue specificity:Placenta and lung.,
Background	This gene encodes a receptor for prostaglandin E2, a metabolite of arachidonic acid which has different biologic activities in a wide range of tissues. Mutations in this gene are associated with aspirin-induced susceptibility to asthma. [provided by RefSeq, Oct 2009],



matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western Blot analysis of various cells using EP2 Monoclonal Antibody

