



PI 3-Kinase p110 γ Monoclonal Antibody


Catalog No	YP-mAb-14914
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	PIK3CG
Immunogen	The antiserum was produced against synthesized peptide derived from human PIK3CG. AA range:881-930
Specificity	PI 3-Kinase p110 γ Monoclonal Antibody detects endogenous levels of PI 3-Kinase p110 γ protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	1 phosphatidylinositol 3 kinase;5-bisphosphate 3-kinase 11 kDa catalytic subunit gamma;5-bisphosphate 3-kinase catalytic subunit gamma isoform;p11 gamma;p11gamma;p12 PI3K;p12-PI3K;Phosphatidylinositol 3 kinase catalytic 11 kD gamma;Phosphatidylinositol 3 kinase gamma;Phosphatidylinositol 3 kinase; catalytic;gamma polypeptide;Phosphatidylinositol 4 5 bisphosphate 3 kinase 11 kDa catalytic subunit gamma;Phosphatidylinositol 4 5 bisphosphate 3 kinase catalytic subunit gamma;Phosphatidylinositol 4 5 bisphosphate 3 kinase catalytic subunit gamma isoform;Phosphatidylinositol-4;Phosphoinositide 3 kinase catalytic gamma polypeptide;Phosphoinositide 3 kinase gamma catalytic subunit; PI3 kinase p11 subunit gamma;PI3-kinase subunit gamma;PI3CG;PI3K;PI3K-gamma;PI3Kgamma;PIK3;Pik3cg;PK3CG_HUMAN;PtdIns-3-kinase subunit gamma;PtdIns-3-kinase subunit p11-gamma;Serine/threonine protein kinase PIK3CG;PI 3-Kinase p110 γ ;Phosphatidylinositol 4;5-bisphosphate 3-kinase 110 kDa catalytic subunit
Observed Band	120kD
Calculated Molecular Weight	120kD
Cell Pathway	Cytoplasm . Cell membrane .



Tissue Specificity	Pancreas, skeletal muscle, liver and heart.
Function	catalytic activity:ATP + 1-phosphatidyl-1D-myo-inositol 4,5-bisphosphate = ADP + 1-phosphatidyl-1D-myo-inositol 3,4,5-trisphosphate.,enzyme regulation:Activated by both the alpha and the beta-gamma G proteins.,function:3-phosphorylates the cellular phosphoinositide PtdIns-4,5-biphosphate (PtdIns(4,5)P2) to produce PtdIns-3, 4,5-triiphosphate (PtdIns(3,4,5)P3). Links G-protein coupled receptor activation to the secondary messenger PtdIns(3,4,5)P3 production.,pathway:Phospholipid metabolism; phosphatidylinositol phosphate biosynthesis.,similarity:Belongs to the PI3/PI4-kinase family.,similarity:Contains 1 PI3K/PI4K domain.,subunit:Heterodimer of a catalytic subunit (PIK3CG/p120) and a regulatory (PIK3R5a/p101) subunit.,tissue specificity:Pancreas, skeletal muscle, liver and heart.,
Background	Phosphoinositide 3-kinases (PI3Ks) phosphorylate inositol lipids and are involved in the immune response. The protein encoded by this gene is a class I catalytic subunit of PI3K. Like other class I catalytic subunits (p110-alpha p110-beta, and p110-delta), the encoded protein binds a p85 regulatory subunit to form PI3K. This gene is located in a commonly deleted segment of chromosome 7 previously identified in myeloid leukemias. Several transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Jun 2015],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis of various cells using PI
3-Kinase p110 γ Monoclonal Antibody