



CATB (Cleaved-Asn126) mouse mAb

Catalog No	YP-mAb-02303
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CTSB CPSB
Immunogen	Synthesized peptide derived from human CATB (Cleaved-Asn126)
Specificity	This antibody detects endogenous levels of Human CATB (Cleaved-Asn126, protein was cleaved amino acid sequence between 126-127)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CATB;CTSB CPSB;CATB (Cleaved-Asn126);Cathepsin B (EC 3.4.22.1;APP secretase;APPS;Cathepsin B1) [Cleaved into: Cathepsin B light chain;Cathepsin B heavy chain]
Observed Band	15 37kD
Calculated Molecular Weight	15 37kD
Cell Pathway	Lysosome . Melanosome . Secreted, extracellular space . Apical cell membrane ; Peripheral membrane protein ; Extracellular side . Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065). Localizes to the lumen of thyroid follicles and to the apical membrane of thyroid epithelial cells (By similarity). .
Tissue Specificity	Expressed in the stratum spinosum of the epidermis. Weak expression is detected in the stratum granulosum.
Function	proteolysis, response to wounding, regulation of cell death, regulation of apoptosis, regulation of programmed cell death,
Background	catalytic activity:Hydrolysis of proteins with broad specificity for peptide bonds. Preferentially cleaves -Arg-Arg- -Xaa bonds in small molecule substrates (thus differing from cathepsin L). In addition to being an endopeptidase, shows



peptidyl-dipeptidase activity, liberating C-terminal dipeptides.,function:Thiol protease which is believed to participate in intracellular degradation and turnover of proteins. Has also been implicated in tumor invasion and metastasis., similarity:Belongs to the peptidase C1 family.,subcellular location:Identified by mass spectrometry in melanosome fractions from stage I to stage IV., subunit:Dimer of a heavy chain and a light chain cross-linked by a disulfide bond.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images