



PSG3 Polyclonal Antibody

Catalog No	YP-Ab-13959
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	IHC;IF;ELISA
Gene Name	PSG3
Immunogen	Synthesized peptide derived from the Internal region of human PSG3.
Specificity	PSG3 Polyclonal Antibody detects endogenous levels of PSG3 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ,IHC: 1/100 - 1/300. ELISA: 1/20000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Carcinoembryonic antigen SG5;Pregnancy-specific beta-1-glycoprotein 3; Pregnancy-specific glycoprotein 3;PS beta G 3;PS-beta-G-3;PSG3;PSBG-3
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Secreted .
Tissue Specificity	Colon,Placenta,Testis,
Function	developmental stage:PSBG are produced in high quantity during pregnancy., similarity:Belongs to the immunoglobulin superfamily. CEA family., similarity:Contains 1 Ig-like V-type (immunoglobulin-like) domain., similarity:Contains 3 Ig-like C2-type (immunoglobulin-like) domains.,
Background	The human pregnancy-specific glycoproteins (PSGs) are a family of proteins that are synthesized in large amounts by placental trophoblasts and released into the maternal circulation during pregnancy. Molecular cloning and analysis of several PSG genes has indicated that the PSGs form a subgroup of the carcinoembryonic antigen (CEA) gene family, which belongs to the immunoglobulin superfamily of genes. Members of the CEA family consist of a single N domain, with structural similarity to the immunoglobulin variable



domains, followed by a variable number of immunoglobulin constant-like A and/or B domains. Most PSGs have an arg-gly-asp (RGD) motif, which has been shown to function as an adhesion recognition signal for several integrins, in the N-terminal domain (summary by Teglund et al., 1994 [PubMed 7851896]). For additional general information about the PSG gene family, see PSG1 (MIM


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).