



RIR2 mouse mAb

Catalog No	YP-mAb-10123
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	RRM2 RR2
Immunogen	Synthesized peptide derived from human RIR2 AA range: 260-310
Specificity	This antibody detects endogenous levels of human RIR2
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, mouse,IgG
Purification	The antibody was affinity-purified from mouse serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	cb111;chunp6884;etID309896.20;R2;reductase M2 polypeptide variant 1; reductase M2 polypeptide variant 2;reductase M2 polypeptide variant 3a; reductase M2 polypeptide variant 3b;reductase M2 polypeptide variant 3c; reductase M2 polypeptide variant 3d;Ribonucleoside-diphosphate reductase subunit M2;Ribonucleotide reductase M2;Ribonucleotide reductase M2 polypeptide;Ribonucleotide reductase M2 subunit;ribonucleotide reductase protein r2 class I;Ribonucleotide reductase regulatory subunit M2; Ribonucleotide reductase small chain;Ribonucleotide reductase small subunit; Ribonucleotide reductase;R2 subunit;RIR2_HUMAN;RR2;RR2M;RRM2;RIR2; RRM2 RR2;Ribonucleoside-diphosphate reductase subunit M2 (EC 1.17.4.1) (Ribonucleotide reductase small chain) (Ribonucleotide reductase small subunit)
Observed Band	45kD
Calculated Molecular Weight	45kDa
Cell Pathway	Cytoplasm . Nucleus . Localized to the cytoplasm in S phase cells. May localize to the nucleus in G2 phase cells. .
Tissue Specificity	
Function	

Background

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images

