



APG7 mouse mAb

Catalog No	YP-mAb-12443
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ATG7 APG7L
Immunogen	Synthesized peptide derived from human APG7 AA range: 521-570
Specificity	This antibody detects endogenous levels of Human,Mouse,Rat APG7
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, mouse,IgG
Purification	The antibody was affinity-purified from mouse serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	GSA7;Autophagy-related protein 7;Autophagy related protein 7;ATG12-activating enzyme E1 ATG7;APG7L;APG7;ATG7 APG7L;Ubiquitin-like modifier-activating enzyme ATG7 (ATG12-activating enzyme E1 ATG7;APG7-like; hAGP7;Ubiquitin-activating enzyme E1-like protein)
Calculated Molecular Weight	78 kDa
Cell Pathway	Cytoplasm . Preautophagosomal structure . Localizes also to discrete punctae along the ciliary axoneme and to the base of the ciliary axoneme. .
Tissue Specificity	Widely expressed, especially in kidney, liver, lymph nodes and bone marrow.
Function	domain:The C-terminal part of the protein is essential for the dimerization and interaction with ATG3 and ATG12.,function:Functions as an E1 enzyme essential for multisubstrates such as GABARAPL1 and ATG12. Forms intermediate conjugates with GABARAPL1 (GABARAPL2, GABARAP or MAP1ALC3). Formation of the final GABARAPL1-PE conjugate is essential for autophagy.,similarity:Belongs to the ATG7 family.,subunit:Homodimer (By similarity). Interacts with ATG3 and ATG12. The complex, composed of ATG3 and ATG7, plays a role in the conjugation of ATG12 to ATG5.,tissue specificity:Widely expressed, especially in kidney, liver, lymph nodes and bone marrow.,

Background

This gene encodes an E1-like activating enzyme that is essential for autophagy and cytoplasmic to vacuole transport. The encoded protein is also thought to modulate p53-dependent cell metabolic stress. cycle pathways during prolonged It has been associated with multiple functions, including axon membrane trafficking, axonal homeostasis, mitophagy, adipose differentiation, and hematopoietic stem cell maintenance. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images