



ARHGAP23 Monoclonal Antibody

Catalog No	YP-mAb-16136
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ARHGAP23
Immunogen	Synthesized peptide derived from ARHGAP23 . at AA range: 470-550
Specificity	ARHGAP23 Monoclonal Antibody detects endogenous levels of ARHGAP23 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	KIAA1501;Rho GTPase Activating Protein 23;Rho GTPase-activating protein 23;Rho-type GTPase-activating protein 23;ARHGAP23
Observed Band	
Calculated Molecular Weight	
Cell Pathway	cytosol,extracellular exosome,
Tissue Specificity	Expressed in placenta, prostate, hippocampus and brain medulla. Also expressed in brain tumor, salivary gland tumor, head and neck tumor.
Function	function:GTPase activator for the Rho-type GTPases by converting them to an inactive GDP-bound state.,similarity:Contains 1 PDZ (DHR) domain.,similarity:Contains 1 PH domain.,similarity:Contains 1 Rho-GAP domain.,tissue specificity:Expressed in placenta, prostate, hippocampus and brain medulla. Also expressed in brain tumor, salivary gland tumor, head and neck tumor.,
Background	The RHO (see ARHA; MIM 165390) family of small GTPases are involved in signal transduction through transmembrane receptors, and they are inactive in the GDP-bound form and active in the GTP-bound form. GTPase-activating proteins, such as ARHGAP23, inactivate RHO family proteins by stimulating



their hydrolysis of GTP (Kato and Kato, 2004 [PubMed 15254754]).[supplied by OMIM, Mar 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images