



ATP5G2 Monoclonal Antibody

Catalog No	YP-mAb-16388
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	ATP5G2
Immunogen	The antiserum was produced against synthesized peptide derived from human ATP5G2. AA range:1-50
Specificity	ATP5G2 Monoclonal Antibody detects endogenous levels of ATP5G2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ATP synthase proteolipid P2;ATP5G2;ATPase protein 9;ATPase subunit c;ATP synthase lipid-binding protein mitochondrial;PSEC0033;ATP synthase lipid-binding protein;mitochondrial
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Mitochondrion membrane; Multi-pass membrane protein.
Tissue Specificity	Liver,Lung,
Function	disease:This protein is the major protein stored in the storage bodies of animals or humans affected with ceroid lipofuscinosis (Batten disease)., function:Mitochondrial membrane ATP synthase (F(1)F(0) ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F(1) - containing the extramembraneous catalytic core and F(0) - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Part



of the complex F(0) domain. A homomeric c-ring of probably 10 subunits is part of the complex rotary element.,miscellaneous:There are three gene

Background

This gene encodes a subunit of mitochondrial ATP synthase. Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. ATP synthase is composed of two linked multi-subunit complexes: the soluble catalytic core, F1, and the membrane-spanning component, Fo, comprising the proton channel. The catalytic portion of mitochondrial ATP synthase consists of 5 different subunits (alpha, beta, gamma, delta, and epsilon) assembled with a stoichiometry of 3 alpha, 3 beta, and single representatives of the gamma, delta, and epsilon subunits. The proton channel likely has nine subunits (a, b, c, d, e, f, g, F6 and 8). There are three separate genes which encode subunit c of the proton channel and they specify precursors with different import sequences but identical

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images