



Na⁺ CP type VII α Monoclonal Antibody

Catalog No	YP-mAb-16474
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SCN7A
Immunogen	The antiserum was produced against synthesized peptide derived from human SCN7A. AA range:771-820
Specificity	Na ⁺ CP type VII α Monoclonal Antibody detects endogenous levels of Na ⁺ CP type VII α protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Alpha 1 type VII collagen;CO7A1_HUMAN;Col7a1;Collagen alpha 1(VII) chain; Collagen alpha-1(VII) chain;Collagen type VII alpha 1;Collagen VII alpha 1 polypeptide;CollagenVII;EBD 1;EBD1;EBDCT;EBR 1;EBR1;LC collagen;Long chain collagen;Long-chain collagen;Na ⁺ CP type VII α ;SCN7A;Sodium channel protein type 7 subunit alpha;SCN6A;Putative voltage-gated sodium channel subunit alpha Nax;Sodium channel protein cardiac and skeletal muscle subunit alpha;Sodium channel protein type VII subunit alpha
Observed Band	200kD
Calculated Molecular Weight	200kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein .
Tissue Specificity	Heart and uterus.
Function	domain:The sequence contains 4 internal repeats, each with 5 hydrophobic segments (S1,S2,S3,S5,S6) and one positively charged segment (S4). Segments S4 are probably the voltage-sensors and are characterized by a series of positively charged amino acids at every third position., function:Mediates the voltage-dependent sodium ion permeability of excitable



membranes. Assuming opened or closed conformations in response to the voltage difference across the membrane, the protein forms a sodium-selective channel through which Na(+) ions may pass in accordance with their electrochemical gradient.,similarity:Belongs to the sodium channel family.,tissue specificity:Heart and uterus.,

Background

This gene encodes one of the many voltage-gated sodium channel proteins. For proper functioning of neurons and muscles during action potentials, voltage-gated sodium channels direct sodium ion diffusion for membrane depolarization. This sodium channel protein has some atypical characteristics; the similarity between the human and mouse proteins is lower compared to other orthologous sodium channel pairs. Also, the S4 segments, which sense voltage changes, have fewer positive charged residues than in other sodium channels; domain 4 has fewer arginine and lysine residues compared to other sodium channel proteins. Several alternatively spliced transcript variants exist, but the full-length natures of all of them remain unknown. [provided by RefSeq, Dec 2011],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images