



AF-4 Monoclonal Antibody

Catalog No	YP-mAb-16926
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	AFF1
Immunogen	The antiserum was produced against synthesized peptide derived from human AF4. AA range:1-50
Specificity	AF-4 Monoclonal Antibody detects endogenous levels of AF-4 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-1:2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	AF4/FMR2 family member 4;AF5Q31;Alf4;ALL1 fused gene from chromosome 5q31;HSPC92;Laf4;Lymphoid nuclear protein related to AF4 like;Major CDK9 elongation factor associated protein;MCEF;MGC17469;RP23-226J16.7;AF-4;AFF1;AF4/FMR2 family member 1;AF4;FEL;MLLT2;PBM1;ALL1-fused gene from chromosome 4 protein;Protein AF-4;Protein FEL;Proto-oncogene AF4
Observed Band	130kD
Calculated Molecular Weight	130kD
Cell Pathway	Nucleus .
Tissue Specificity	Epithelium,PCR rescued clones,Placenta,
Function	disease:A chromosomal aberration involving AFF1 is associated with acute leukemias. Translocation t(4;11)(q21;q23) with MLL/HRX. The result is a rogue activator protein.,similarity:Belongs to the AF4 family.,
Background	This gene encodes a member of the AF4/ lymphoid nuclear protein related to AF4/Fragile X E mental retardation syndrome family of proteins, which have been implicated in childhood lymphoblastic leukemia, Fragile X E site mental retardation, and ataxia. It is the prevalent mixed-lineage leukemia fusion gene



associated with spontaneous acute lymphoblastic leukemia. Members of this family have three conserved domains: an N-terminal homology domain, an AF4/lymphoid nuclear protein related to AF4/Fragile X E mental retardation syndrome domain, and a C-terminal homology domain. The protein functions as a regulator of RNA polymerase II-mediated transcription through elongation and chromatin remodeling functions. Through RNA interference screens, this gene has been shown to promote the expression of CD133, a plasma membrane glycoprotein required for leukemia cell survival. Alternative splicing results in mu

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images