



# NOB1 mouse mAb

|                                    |                                                                                                                                                                          |
|------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Catalog No</b>                  | YP-mAb-18265                                                                                                                                                             |
| <b>Isotype</b>                     | IgG                                                                                                                                                                      |
| <b>Reactivity</b>                  | Human;Mouse;Rat                                                                                                                                                          |
| <b>Applications</b>                | WB                                                                                                                                                                       |
| <b>Gene Name</b>                   | NOB1 ART4 NOB1P PSMD8BP1 MSTP158                                                                                                                                         |
| <b>Immunogen</b>                   | Synthesized peptide derived from human NOB1                                                                                                                              |
| <b>Specificity</b>                 | This antibody detects endogenous levels of NOB1 at Human, Mouse,Rat                                                                                                      |
| <b>Formulation</b>                 | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                                                                                                  |
| <b>Source</b>                      | Monoclonal, Mouse,IgG                                                                                                                                                    |
| <b>Purification</b>                | The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.                                                     |
| <b>Dilution</b>                    | WB 1:500-1:2000                                                                                                                                                          |
| <b>Concentration</b>               | 1 mg/ml                                                                                                                                                                  |
| <b>Purity</b>                      | ≥90%                                                                                                                                                                     |
| <b>Storage Stability</b>           | -20°C/1 year                                                                                                                                                             |
| <b>Synonyms</b>                    | NOB1P;MSTP158;MST158;EC:3.1.-.-;ART4;NOB1;NOB1 ART4 NOB1P PSMD8BP1 MSTP158;RNA-binding protein NOB1 (Phosphorylation regulatory protein HP-10) (Protein ART-4)           |
| <b>Observed Band</b>               | 47-50 kDa                                                                                                                                                                |
| <b>Calculated Molecular Weight</b> | 47 kDa                                                                                                                                                                   |
| <b>Cell Pathway</b>                | Nucleus .                                                                                                                                                                |
| <b>Tissue Specificity</b>          | Detected in liver, lung, placenta, endothelial cells and spleen.                                                                                                         |
| <b>Function</b>                    | May play a role in mRNA degradation (Probable). Endonuclease required for processing of 20S pre-rRNA precursor and biogenesis of 40S ribosomal subunits (By similarity). |
| <b>Background</b>                  |                                                                                                                                                                          |
| <b>matters needing attention</b>   | Avoid repeated freezing and thawing!                                                                                                                                     |
| <b>Usage suggestions</b>           | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.                                        |



## Products Images