



VPS24 Mouse mAb

Catalog No	YP-mAb-19159
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	CHMP3 CGI149 NEDF VPS24 CGI-149
Immunogen	Synthesized peptide derived from human VPS24
Specificity	This antibody detects endogenous levels of VPS24 at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal,Mouse,IgG
Purification	The antibody was affinity-purified from mouse antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	NEDF;VPS24;CGI-149;CHMP3 CGI149 NEDF VPS24 CGI-149;Charged multivesicular body protein 3 (Chromatin-modifying protein 3) (Neuroendocrine differentiation factor) (Vacuolar protein sorting-associated protein 24) (hVps24)
Observed Band	29kDa
Calculated Molecular Weight	25kDa
Cell Pathway	Cytoplasm, cytosol. Membrane; Lipid-anchor. Endosome. Late endosome membrane . Localizes to the midbody of dividing cells.
Tissue Specificity	Widely expressed. Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas.
Function	Probable core component of the endosomal sorting required for transport complex III (ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. The MVB pathway appears to require the sequential function of ESCRT-O, -I,-II and -III complexes. ESCRT-III proteins mostly dissociate from the invaginating membrane before the ILV is released.



The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis and the budding of enveloped viruses (HIV-1 and other lentiviruses). ESCRT-III proteins are believed to mediate the necessary vesicle extrusion and/or membrane fission activities, possibly in conjunction with the AAA ATPase VPS4. Selectively binds to phosphatidylinositol 3,5-bisphosphate PtdIns(3,5)P2 and PtdIns(3,4)P2 in preference to other phosphoinositides tested. Involved in late stages of cytokinesis. Plays a role in endosomal sorting/trafficking of EGF receptor. Isoform 2 prevents stress-mediated cell death and accumulation of reactive oxygen species when expressed in yeast cells.

Background**matters needing attention**

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images