



MEK-6 Monoclonal Antibody

Catalog No	YP-Ab-14175
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;IF;FCM;ELISA
Gene Name	MAP2K6
Immunogen	Purified recombinant fragment of human MEK-6 expressed in E. Coli.
Specificity	MEK-6 Monoclonal Antibody detects endogenous levels of MEK-6 protein.
Formulation	Ascitic fluid containing 0.03% sodium azide,0.5% BSA, 50%glycerol.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. Flow cytometry: 1/200 - 1/400. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MKK6 (Ser207);Phospho-MEK6;Phospho-MEK6(Ser207);Phospho-MKK3 (Ser189);Phospho-MKK3;MEK-6;MAP2K6;Dual specificity mitogen-activated protein kinase kinase 6;MEK6;MKK6;PRKMK6;SKK3;MAP kinase kinase 6; MAPKK 6;MAPK/ERK kinase 6;MEK 6;Stress-activated protein kinase kinase 3; SAPK kinase 3;SAPKK-3;SAPKK3
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Nucleus . Cytoplasm . Cytoplasm, cytoskeleton . Binds to microtubules.
Tissue Specificity	Isoform 2 is only expressed in skeletal muscle. Isoform 1 is expressed in skeletal muscle, heart, and in lesser extent in liver or pancreas.
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein.,enzyme regulation:Probably activated by dual phosphorylation on Ser-207 and Thr-211., function:Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in MAP kinase p38 exclusively.,induction:Strongly activated by UV, anisomycin, and osmotic shock but not by phorbol esters, NGF or EGF., PTM:Acetylation of Ser-207 and Thr-211 by Yersinia yopJ prevents phosphorylation and activation, thus blocking the MAPK signaling pathway., PTM:Weakly autophosphorylated.,similarity:Belongs to the protein kinase



superfamily.,similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. MAP kinase kinase subfamily.,similarity:Contains 1 protein kinase domain.,subunit:Interacts with Yersinia yopJ.,tissue specificity:Isoform 2 is only expressed in skeletal muscle. Isoform 1, on the other hand, is found

Background

This gene encodes a member of the dual specificity protein kinase family, which functions as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This protein phosphorylates and activates p38 MAP kinase in response to inflammatory cytokines or environmental stress. As an essential component of p38 MAP kinase mediated signal transduction pathway, this gene is involved in many cellular processes such as stress induced cell cycle arrest, transcription activation and apoptosis. [provided by RefSeq, Jul 2008],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western Blot analysis using MEK-6 Monoclonal Antibody against HepG2 (1), MCF-7 (2) and NIH/3T3 (3) cell lysate.



Immunofluorescence analysis of HeLa cells using MEK-6 Monoclonal Antibody (green). Blue: DRAQ5 fluorescent DNA dye.



Flow cytometric analysis of HeLa cells using MEK-6 Monoclonal Antibody (green) and negative control (purple).

