



MKP-1 (phospho Ser359) Polyclonal Antibody

Catalog No	YP-Ab-14534
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	IHC;IF;ELISA
Gene Name	DUSP1
Immunogen	The antiserum was produced against synthesized peptide derived from human MKP1 around the phosphorylation site of Ser359. AA range:318-367
Specificity	Phospho-MKP-1 (S359) Polyclonal Antibody detects endogenous levels of MKP-1 protein only when phosphorylated at S359.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/10000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Dual specificity protein phosphatase 12;EC:3.1.3.16;MKP-1;DUSP1;Dual specificity protein phosphatase 1;CL100;MKP1;PTPN10;VH1;Dual specificity protein phosphatase hVH1;Mitogen-activated protein kinase phosphatase 1; MAP kinase phosphatase 1;Protein-tyrosine phosphatase CL100
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Nucleus .
Tissue Specificity	Expressed at high levels in the lung, liver placenta and pancreas. Moderate levels seen in the heart and skeletal muscle. Lower levels found in the brain and kidney.
Function	catalytic activity:A phosphoprotein + H(2)O = a protein + phosphate.,catalytic activity:Protein tyrosine phosphate + H(2)O = protein tyrosine + phosphate., function: Dual specificity phosphatase that dephosphorylates MAP kinase ERK2 on both 'Thr-183' and 'Tyr-185'. ,induction: By oxidative stress and heat shock., similarity: Belongs to the protein-tyrosine phosphatase family. Non-receptor class dual specificity subfamily.,similarity: Contains 1 rhodanese domain., similarity: Contains 1 tyrosine-protein phosphatase domain.,tissue



specificity: Expressed at high levels in the lung, liver placenta and pancreas. Moderate levels seen in the heart and skeletal muscle. Lower levels found in the brain and kidney.,

Background

The expression of DUSP1 gene is induced in human skin fibroblasts by oxidative/heat stress and growth factors. It specifies a protein with structural features similar to members of the non-receptor-type protein-tyrosine phosphatase family, and which has significant amino-acid sequence similarity to a Tyr/Ser-protein phosphatase encoded by the late gene H1 of vaccinia virus. The bacterially expressed and purified DUSP1 protein has intrinsic phosphatase activity, and specifically inactivates mitogen-activated protein (MAP) kinase in vitro by the concomitant dephosphorylation of both its phosphothreonine and phosphotyrosine residues. Furthermore, it suppresses the activation of MAP kinase by oncogenic ras in extracts of *Xenopus* oocytes. Thus, DUSP1 may play an important role in the human cellular response to environmental stress as well as in the negative regulation of cellular proliferation.

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using MKP1 (Phospho-Ser359) Antibody



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma, using MKP1 (Phospho-Ser359) Antibody. The picture on the right is blocked with the phospho peptide.