



MLK1 Polyclonal Antibody

Catalog No	YP-Ab-14853
Isotype	IgG
Reactivity	Human;Mouse
Applications	IHC;IF;ELISA
Gene Name	MAP3K9
Immunogen	The antiserum was produced against synthesized peptide derived from human MAP3K9. AA range:561-610
Specificity	MLK1 Polyclonal Antibody detects endogenous levels of MLK1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/20000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MEKK1;MEKK9;Mitogen activated protein kinase kinase kinase 1;Mitogen activated protein kinase kinase kinase 9;Mixed lineage kinase 1 (tyr and ser/thr specificity);Mixed lineage kinase 2;MKN28 derived nonreceptor_type serine/threonine kinase;MKN28 kinase;MLK1;MLK2;MST;PRKE1;Protein kinase MST; MAP3K9;Mitogen-activated protein kinase kinase kinase 9;Mixed lineage kinase 1
Observed Band	
Calculated Molecular Weight	
Cell Pathway	intracellular,integral component of membrane,
Tissue Specificity	Expressed in epithelial tumor cell lines of colonic, breast and esophageal origin.
Function	catalytic activity:ATP + a protein = ADP + a phosphoprotein., cofactor:Magnesium.,enzyme regulation:Homodimerization via the leucine zipper domains is required for autophosphorylation and subsequent activation., function:Activates the JUN N-terminal pathway.,PTM:Autophosphorylation on serine and threonine residues within the activation loop plays a role in enzyme activation. Thr-312 is likely to be the main autophosphorylation site., similarity:Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase



family. MAP kinase kinase kinase subfamily.,similarity:Contains 1 protein kinase domain.,similarity:Contains 1 SH3 domain.,subunit:Homodimer.,tissue specificity:Expressed in epithelial tumor cell lines of colonic, breast and esophageal origin.,

Background

MAP3K9 (Mitogen-Activated Protein Kinase Kinase 9) is a Protein Coding gene. Diseases associated with MAP3K9 include retroperitoneal neuroblastoma. Among its related pathways are MAP Kinase Signaling and TGF-Beta Pathway. GO annotations related to this gene include protein homodimerization activity and protein kinase activity. An important paralog of this gene is KSR1.


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using MAP3K9 Antibody. The picture on the right is blocked with the synthesized peptide.