



PP2A-B55-β Polyclonal Antibody

Catalog No	YP-Ab-14948
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC
Gene Name	PPP2R2B
Immunogen	Synthesized peptide derived from PP2A-B55-β . at AA range: 90-170
Specificity	PP2A-B55-β Polyclonal Antibody detects endogenous levels of PP2A-B55-β protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	PP2A-B55-β;PPP2R2B;Serine/threonine-protein phosphatase 2A 55 kDa regulatory subunit B beta isoform;PP2A subunit B isoform B55-beta;PP2A subunit B isoform PR55-beta;PP2A subunit B isoform R2-beta;PP2A subunit B isoform beta
Observed Band	51kD
Calculated Molecular Weight	51kD
Cell Pathway	[Isoform 1]: Cytoplasm . Cytoplasm, cytoskeleton . Membrane .; [Isoform 2]: Cytoplasm . Mitochondrion . Mitochondrion outer membrane . Under basal conditions, localizes to both cytosolic and mitochondrial compartments. Relocalizes from the cytosolic to the mitochondrial compartment during apoptosis. Its targeting to the outer mitochondrial membrane (OMM) involves an association with import receptors of the TOM complex and is required to promote proapoptotic activity (By similarity). .
Tissue Specificity	Brain.
Function	disease:Defects in PPP2R2B are the cause of spinocerebellar ataxia type 12 (SCA12) [MIM:604326]. Spinocerebellar ataxia is a clinically and genetically heterogeneous group of cerebellar disorders. Patients show progressive



incoordination of gait and often poor coordination of hands, speech and eye movements, due to degeneration of the cerebellum with variable involvement of the brainstem and spinal cord. SCA12 is an autosomal dominant cerebellar ataxia (ADCA).,function:The B regulatory subunit might modulate substrate selectivity and catalytic activity, and also might direct the localization of the catalytic enzyme to a particular subcellular compartment.,similarity:Belongs to the phosphatase 2A regulatory subunit B family.,similarity:Contains 7 WD repeats.,subunit:PP2A consists of a common heterodimeric core enzyme, composed of a 36 kDa catalytic subunit (subunit C) and a 65 kDa constan

Background

protein phosphatase 2 regulatory subunit Bbeta(PPP2R2B) Homo sapiens The product of this gene belongs to the phosphatase 2 regulatory subunit B family. Protein phosphatase 2 is one of the four major Ser/Thr phosphatases, and it is implicated in the negative control of cell growth and division. It consists of a common heteromeric core enzyme, which is composed of a catalytic subunit and a constant regulatory subunit, that associates with a variety of regulatory subunits. The B regulatory subunit might modulate substrate selectivity and catalytic activity. This gene encodes a beta isoform of the regulatory subunit B55 subfamily. Defects in this gene cause autosomal dominant spinocerebellar ataxia 12 (SCA12), a disease caused by degeneration of the cerebellum, sometimes involving the brainstem and spinal cord, and in resulting in poor coordination of speech and body movements. Multiple alternatively spliced variants, which encode different isoform


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human cervical carcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).