



RERG Polyclonal Antibody

Catalog No	YP-Ab-16235
Isotype	IgG
Reactivity	Human;Mouse
Applications	IHC;IF;ELISA
Gene Name	RERG
Immunogen	Synthesized peptide derived from RERG . at AA range: 70-150
Specificity	RERG Polyclonal Antibody detects endogenous levels of RERG protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC: 1/100 - 1/300. ELISA: 1/5000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	EC:3.6.5.2;Ras-related and estrogen-regulated growth inhibitor;RERC;RERG
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Cytoplasm .
Tissue Specificity	Detected in heart, brain, placenta, lung, liver, skin, kidney and pancreas. Detected in estrogen receptor-positive breast-derived cell lines, but not in estrogen receptor-negative cell lines. Expression is decreased or lost in a significant proportion of primary breast tumors with poor clinical prognosis.
Function	disease:Loss of RERG expression may contribute to tumorigenesis in breast. RERG expression is decreased or lost in a significant proportion of primary breast tumors with poor clinical prognosis. High expression of RERG correlates with long relapse-free survival times.,function:Binds GDP/GTP and possesses intrinsic GTPase activity. Has higher affinity for GDP than for GTP. In cell lines overexpression leads to a reduction in the rate of proliferation, colony formation and in tumorigenic potential.,induction:Up-regulated by estradiol. Down-regulated by tamoxifen.,similarity:Belongs to the small GTPase superfamily. Ras family.,tissue specificity:Detected in heart, brain, placenta, lung, liver, skin, kidney and pancreas. Detected in estrogen receptor-positive breast-derived cell



lines, but not in estrogen receptor-negative cell lines.,

Background

RERG, a member of the RAS superfamily of GTPases, inhibits cell proliferation and tumor formation (Finlin et al., 2001 [PubMed 11533059]).[supplied by OMIM, Mar 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Tris-EDTA,pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200(4° overnight.3,Secondary antibody was diluted at 1:200(room temperature, 45min).