



Kv10.1 Polyclonal Antibody


Catalog No	YP-Ab-16313
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	IHC;IF
Gene Name	KCNG3
Immunogen	Synthetic Peptide of Kv10.1 AA range: 335-385
Specificity	Kv10.1 protein(A259) detects endogenous levels of Kv10.1
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using specific immunogen.
Dilution	IHC 1:100-200. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	EAG;EAG channel 1;EAG1;Ether a go go potassium channel 1;Ether a go go; Drosophila;homolog of;Ether-a-go-go potassium channel 1;ether-a-go-go;h eag; h-eag;hEAG1;Kcnh1;KCNH1_HUMAN;Kv1.1;M eag;MGC124419;MGC12442; MGC142269;Potassium voltage gated channel subfamily H (eag related); member 1;Potassium voltage-gated channel subfamily H member 1;Voltage gated potassium channel subunit Kv1.1;Voltage-gated potassium channel subunit Kv1.1;Kv10.1;KCNG3;Potassium voltage-gated channel subfamily G member 3 (Voltage-gated potassium channel subunit Kv10.1) (Voltage-gated potassium channel subunit Kv6.3);Potassium voltage-gated channel subfamily G member 3;Voltage-gated potassium channel subunit Kv10.1;Voltage-gated potassium channel subunit Kv6.3
Observed Band	50kD
Calculated Molecular Weight	50kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein . Cytoplasm . Has to be associated with KCNB1 or possibly another partner to get inserted in the plasma membrane (PubMed:12060745). Colocalizes with KCNB1 at the plasma membrane (PubMed:12060745, PubMed:19074135). Remains intracellular in the absence of KCNB1 (PubMed:12060745). .



Tissue Specificity	Expressed in the brain, liver, testis, small intestine, colon, thymus and adrenal gland (PubMed:11852086, PubMed:12060745).
Function	domain:The segment S4 is probably the voltage-sensor and is characterized by a series of positively charged amino acids at every third position., function:Potassium channel subunit. Modulates channel activity., similarity:Belongs to the potassium channel family. G subfamily.,subcellular location:Has to be associated with KCNB1 or possibly another partner to get inserted in the plasma membrane. Remains intracellular in the absence of KCNB1.,subunit:Heteromultimer with KCNB1, KCNC1 and KCNF1. Does not form homomultimers.,tissue specificity:Detected in many parts of the brain with the exception of the cerebellum, in testis, pancreas, lung, kidney, ovary, small intestine, colon, thymus, adrenal gland and spinal cord.,
Background	Voltage-gated potassium (Kv) channels represent the most complex class of voltage-gated ion channels from both functional and structural standpoints. Their diverse functions include regulating neurotransmitter release, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smooth muscle contraction, and cell volume. This gene encodes a member of the potassium channel, voltage-gated, subfamily G. This member is a gamma subunit functioning as a modulatory molecule. Alternative splicing results in two transcript variants encoding distinct isoforms. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Immunohistochemical analysis of paraffin-embedded Rat Brain Tissue using Kv10.1 Rabbit pAb diluted at 1:200.



Immunohistochemical analysis of paraffin-embedded Mouse Brain Tissue using Kv10.1 Rabbit pAb diluted at 1:200.