



GABAA R β 1 Polyclonal Antibody

Catalog No	YP-Ab-16417
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	IHC;IF;WB;ELISA
Gene Name	GABRB1
Immunogen	The antiserum was produced against synthesized peptide derived from human GABA-RB. AA range:401-450
Specificity	GABAA R β 1 Polyclonal Antibody detects endogenous levels of GABAA R β 1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 IHC: 1/100 - 1/300. ELISA: 1/40000.. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	ECA5;GABA alpha receptor beta-2 subunit;GABA(A) receptor subunit beta-3; GABAA receptor beta 3 subunit;GABAA receptor subunit beta 3;GABR B3; Gabrb3;Gamma aminobutyric acid (GABA) A receptor beta 3;Gamma aminobutyric acid receptor subunit beta 3;Gamma-aminobutyric acid receptor subunit beta-3;GBRB3_HUMAN;MGC951;GABAA R β 1;GABRB1;Gamma-aminobutyric acid receptor subunit beta-1;GABA(A) receptor subunit beta-1
Observed Band	55kD
Calculated Molecular Weight	55kD
Cell Pathway	Cell junction, synapse, postsynaptic cell membrane ; Multi-pass membrane protein . Cell membrane ; Multi-pass membrane protein .
Tissue Specificity	Brain,Epithelium,
Function	function:GABA, the major inhibitory neurotransmitter in the vertebrate brain, mediates neuronal inhibition by binding to the GABA/benzodiazepine receptor and opening an integral chloride channel.,online information:Forbidden fruit - Issue 56 of March 2005,similarity:Belongs to the ligand-gated ionic channel (TC 1.A.9) family.,subunit:Binds UBQLN1 (By similarity). Generally pentameric. There are five types of GABA(A) receptor chains: alpha, beta, gamma, delta,



and rho.,

Background

The gamma-aminobutyric acid (GABA) A receptor is a multisubunit chloride channel that mediates the fastest inhibitory synaptic transmission in the central nervous system. This gene encodes GABA A receptor, beta 1 subunit. It is mapped to chromosome 4p12 in a cluster comprised of genes encoding alpha 4, alpha 2 and gamma 1 subunits of the GABA A receptor. Alteration of this gene is implicated in the pathogenetics of schizophrenia. [provided by RefSeq, Jul 2008],



matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using GABA-RB Antibody. The picture on the right is blocked with the synthesized peptide.