



KV9.2 Polyclonal Antibody

Catalog No	YP-Ab-16459
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC
Gene Name	KCNS2
Immunogen	The antiserum was produced against synthesized peptide derived from human KCNS2. AA range:197-246
Specificity	KV9.2 Polyclonal Antibody detects endogenous levels of KV9.2 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	KCNS2;KIAA1144;KV9.2;Potassium voltage-gated channel subfamily S member 2;Delayed-rectifier K(+) channel alpha subunit 2;Voltage-gated potassium channel subunit Kv9.2
Observed Band	54kD
Calculated Molecular Weight	54kD
Cell Pathway	Cell membrane ; Multi-pass membrane protein . May not reach the plasma membrane but remain in an intracellular compartment in the absence of KCNB1 or KCNB2. .
Tissue Specificity	Brain,
Function	domain:The segment S4 is probably the voltage-sensor and is characterized by a series of positively charged amino acids at every third position., function:Potassium channel subunit. Modulates channel activity and reduces the ion flow.,similarity:Belongs to the potassium channel family. S subfamily., subcellular location:May not reach the plasma membrane but remain in an intracellular compartment in the absence of KCNB1.,subunit:Heteromultimer with KCNB1 and with KCNB2. Does not form homomultimers. Might also bind to other channel proteins.,

**Background**

domain:The segment S4 is probably the voltage-sensor and is characterized by a series of positively charged amino acids at every third position.,
function:Potassium channel subunit. Modulates channel activity and reduces the ion flow.,similarity:Belongs to the potassium channel family. S subfamily.,
subcellular location:May not reach the plasma membrane but remain in an intracellular compartment in the absence of KCNB1.,subunit:Heteromultimer with KCNB1 and with KCNB2. Does not form homomultimers. Might also bind to other channel proteins.,

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




Products Images



Western Blot analysis of various cells using KV9.2 Polyclonal Antibody



Western blot analysis of lysates from HepG2 cells, using KCNS2 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).