



ENaC γ Polyclonal Antibody

Catalog No	YP-Ab-16517
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	SCNN1G
Immunogen	The antiserum was produced against synthesized peptide derived from human ENaC gamma. AA range:132-181
Specificity	ENaC γ Polyclonal Antibody detects endogenous levels of ENaC γ protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	Amiloride sensitive epithelial sodium channel gamma subunit;Amiloride sensitive sodium channel subunit gamma;Amiloride-sensitive sodium channel subunit gamma;BESC3;ENaC gamma subunit;ENaCG;ENaCgamma;Epithelial Na(+) channel subunit gamma;Epithelial Na+ channel subunit gamma;Gamma ENaC; Gamma NaCH;Gamma-ENaC;Gamma-NaCH;Nonvoltage gated sodium channel 1 subunit gamma;Nonvoltage-gated sodium channel 1 subunit gamma; PHA 1;PHA1;SCNEG;SCNN 1G;SCNN1G;SCNNG_HUMAN;Sodium channel non voltage gated 1 gamma subunit;Sodium channel nonvoltage gated 1 gamma;ENaC γ
Observed Band	80kD
Calculated Molecular Weight	80kD
Cell Pathway	Apical cell membrane ; Multi-pass membrane protein . Apical membrane of epithelial cells. .
Tissue Specificity	Expressed in kidney (at protein level).
Function	disease:Defects in SCNN1G are a cause of Liddle syndrome [MIM:177200]. It is an autosomal dominant disorder characterized by pseudoaldosteronism and hypertension associated with hypokalemic alkalosis. The disease is caused by



constitutive activation of the renal epithelial sodium channel.,function:Sodium permeable non-voltage-sensitive ion channel inhibited by the diuretic amiloride. Mediates the electrodiffusion of the luminal sodium (and water, which follows osmotically) through the apical membrane of epithelial cells. Controls the reabsorption of sodium in kidney, colon, lung and sweat glands. Also plays a role in taste perception.,PTM:Phosphorylated on serine and threonine residues.,PTM:Ubiquitinated; this targets individual subunits for endocytosis and proteasome-mediated degradation.,similarity:Belongs to the amiloride-sensitive sodium channel family.,subcellular location:Apical me

Background

Nonvoltage-gated, amiloride-sensitive, sodium channels control fluid and electrolyte transport across epithelia in many organs. These channels are heteromeric complexes consisting of 3 subunits: alpha, beta, and gamma. This gene encodes the gamma subunit, and mutations in this gene have been associated with Liddle syndrome. [provided by RefSeq, Apr 2009],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western Blot analysis of extracts from A549 cells, using ENaC γ Polyclonal Antibody. Antibody was diluted at 1:500. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysates from A549 cells, using ENaC γ antibody.