



Cyclin D2 mouse mAb

Catalog No	YP-Ab-16569
Isotype	IgG
Reactivity	Human
Applications	WB
Gene Name	ccnd2
Immunogen	Purified recombinant human Cyclin D2 protein fragments expressed in E.coli.
Specificity	This antibody detects over-expressed levels of Cyclin D2 and does not cross-react with related proteins.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Monoclonal, Mouse
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
Dilution	wb 1:1000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CCND2;G1/S-specific cyclin-D2;G1/S specific cyclin D2;Cyclin D;Cyclin D2;CCND 2;CCND2_HUMAN;CyclinD2;KIAK0002;MGC102758.
Observed Band	38kD
Calculated Molecular Weight	38kD
Cell Pathway	Nucleus . Cytoplasm . Nucleus membrane . Cyclin D-CDK4 complexes accumulate at the nuclear membrane and are then translocated into the nucleus through interaction with KIP/CIP family members. .; [Isoform 2]: Cytoplasm .
Tissue Specificity	Bone marrow,Heart,Placenta,Uterus,
Function	function:Essential for the control of the cell cycle at the G1/S (start) transition., similarity:Belongs to the cyclin family.,similarity:Belongs to the cyclin family. Cyclin D subfamily.,subunit:Interacts with the CDK4 and CDK6 protein kinases to form a serine/threonine kinase holoenzyme complex. The cyclin subunit imparts substrate specificity to the complex.,
Background	The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK



kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with CDK4 or CDK6 and functions as a regulatory subunit of the complex, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with and be involved in the phosphorylation of tumor suppressor protein Rb. Knockout studies of the homologous gene in mouse suggest the essential roles of this gene in ovarian granulosa and germ cell proliferation. High level expression of this gene was observed in ovarian and testicular tumors. Mutations in this gene are associated with megalenceph


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot detection of Cyclin D2 in CHO-K1 cell lysate(A)and CHO-K1 transfected by Cyclin D2-fragment EGFP fusion protein(B)cell lysate using Cyclin D2 mouse mAb (1:1000 diluted).Predicted band size:38KDa.Observed band size:38KDa.