



Cleaved-Integrin α 5 HC (F42) Polyclonal Antibody

Catalog No	YP-Ab-16804
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	ITGA5
Immunogen	The antiserum was produced against synthesized peptide derived from human ITGA5. AA range:23-72
Specificity	Cleaved-Integrin α 5 HC (F42) Polyclonal Antibody detects endogenous levels of fragment of activated Integrin α 5 HC protein resulting from cleavage adjacent to F42.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	$\geq 90\%$
Storage Stability	-20°C/1 year
Synonyms	CD49 antigen-like family member E;CD49e;Fibronectin receptor subunit alpha; Fibronectin receptor;alpha subunit;FNRA;Integrin alpha 5 (fibronectin receptor alpha);Integrin alpha-5;Integrin alpha-5 light chain;Integrin alpha-F;Integrin;alpha 5 (fibronectin receptor;alpha polypeptide);ITA5_HUMAN;Itga5;Very late activation protein 5;VLA-5;VLA5;VLA5A;Integrin α 5;CD antigen CD49e
Observed Band	92kD
Calculated Molecular Weight	92kD
Cell Pathway	Membrane; Single-pass type I membrane protein. Cell junction, focal adhesion . Cell surface .
Tissue Specificity	Liver,Placenta,
Function	function:Integrin alpha-5/beta-1 is a receptor for fibronectin and fibrinogen. It recognizes the sequence R-G-D in its ligands. In case of HIV-1 infection, the interaction with extracellular viral Tat protein seems to enhance angiogenesis in Kaposi's sarcoma lesions.,similarity:Belongs to the integrin alpha chain family., similarity:Contains 7 FG-GAP repeats.,subunit:Heterodimer of an alpha and a



beta subunit. The alpha subunit is composed of an heavy and a light chain linked by a disulfide bond. Alpha-5 associates with beta-1. Interacts with HPS5 and NISCH. Interacts with HIV-1 Tat. Interacts with RAB21.,

Background

integrin subunit alpha 5(ITGA5) Homo sapiens The product of this gene belongs to the integrin alpha chain family. Integrins are heterodimeric integral membrane proteins composed of an alpha subunit and a beta subunit that function in cell surface adhesion and signaling. The encoded preproprotein is proteolytically processed to generate light and heavy chains that comprise the alpha 5 subunit. This subunit associates with the beta 1 subunit to form a fibronectin receptor. This integrin may promote tumor invasion, and higher expression of this gene may be correlated with shorter survival time in lung cancer patients. Note that the integrin alpha 5 and integrin alpha V subunits are encoded by distinct genes. [provided by RefSeq, Oct 2015],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from NIH/3T3 cells, treated with etoposide 25uM 24h, using ITGA5 (heavy chain,Cleaved-Phe42) Antibody. The lane on the right is blocked with the synthesized peptide.