



ODB2 rabbit pAb

Catalog No	YP-Ab-17134
Isotype	IgG
Reactivity	Human, Mouse
Applications	IHC, WB
Gene Name	DBT BCATE2
Immunogen	Synthesized peptide derived from human C-terminal ODB2
Specificity	This antibody detects endogenous levels of ODB2 at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Rabbit, polyclonal
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000 IHC 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	XPE; DDBB; UV-DDB2; DDB2; DBT; DBT BCATE2; Lipoamide acyltransferase component of branched-chain alpha-keto acid dehydrogenase complex; mitochondrial (EC 2.3.1.168) (Branched-chain alpha-keto acid dehydrogenase complex component E2) (BCKAD-E2); mitochondrial (EC 2.3.1.168) (Branched-chain alpha-keto acid dehydrogenase complex component E2) (BCKAD-E2) (BCKADE2) (Dihydrolipoamide acetyltransferase component of branched-chain alpha-keto acid dehydrogenase complex) (Dihydrolipoamide branched chain transacylase) (Dihydrolipoyllysine-residue (2-methylpropanoyl)transferase)
Observed Band	48kDa
Calculated Molecular Weight	48kDa
Cell Pathway	Mitochondrion matrix.
Tissue Specificity	
Function	The branched-chain alpha-keto dehydrogenase complex catalyzes the overall conversion of alpha-keto acids to acyl-CoA and CO ₂ . It contains multiple copies of three enzymatic components: branched-chain alpha-keto acid decarboxylase (E1), lipoamide acyltransferase (E2) and lipoamide dehydrogenase (E3). Within this complex, the catalytic function of this enzyme is to accept, and to transfer to coenzyme A, acyl groups that are generated by the



branched-chain alpha-keto acid decarboxylase component.

Background

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images