



Nucleolin Rabbit pAb

Catalog No	YP-Ab-18964
Isotype	IgG
Reactivity	Human,Mouse,Rat
Applications	WB;ELISA
Gene Name	NCL
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	NUCL Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000; ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	C23;Nsr1;Nucleolin;NCL;Nucleolin (Protein C23)
Observed Band	110kDa
Calculated Molecular Weight	77kDa
Cell Pathway	Nucleus, nucleolus. Cytoplasm. Localized in cytoplasmic mRNP granules containing untranslated mRNAs.
Tissue Specificity	Brain,Cervix carcinoma,Epithelium,Hair follicle dermal papilla,Muscle,Retina,
Function	Nucleolin is the major nucleolar protein of growing eukaryotic cells. It is found associated with intranucleolar chromatin and pre-ribosomal particles. It induces chromatin decondensation by binding to histone H1. It is thought to play a role in pre-rRNA transcription and ribosome assembly. May play a role in the process of transcriptional elongation. Binds RNA oligonucleotides with 5'-UUAGGG-3' repeats more tightly than the telomeric single-stranded DNA 5'-TTAGGG-3' repeats.,sequence Caution:Unlikely isoform. Aberrant splice sites., similarity:Contains 4 RRM (RNA recognition motif) domains.,subunit:Interacts with APTX and NSUN2. Component of the SWAP complex that consists of NPM1, NCL/nucleolin, PARP1 and SWAP70. Component of a complex which is at least composed of HTATSF1/Tat-SF1, the P-TEFb complex components CDK9 and CCNT1, RNA polymerase II, SUPT5H, and NCL/nucleolin.,

**Background**

Nucleolin (NCL), a eukaryotic nucleolar phosphoprotein, is involved in the synthesis and maturation of ribosomes. It is located mainly in dense fibrillar regions of the nucleolus. Human NCL gene consists of 14 exons with 13 introns and spans approximately 11kb. The intron 11 of the NCL gene encodes a small nucleolar RNA, termed U20.

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images