



MEIS2 Rabbit pAb

Catalog No	YP-Ab-19389
Isotype	IgG
Reactivity	Human,Mouse
Applications	WB
Gene Name	MEIS2 MRG1
Immunogen	Synthesized peptide derived from human MEIS2. AA range:88-188
Specificity	This antibody detects endogenous levels of MEIS2 at Human, Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	MEIS2 MRG1;Homeobox protein Meis2 (Meis1-related protein 1)
Observed Band	
Calculated Molecular Weight	52kD
Cell Pathway	Nucleus . Cytoplasm , perinuclear region .
Tissue Specificity	Expressed in various tissues. Expressed at high level in the lymphoid organs of hematopoietic tissues. Also expressed in some regions of the brain , such as the putamen.
Function	Involved in transcriptional regulation. Binds to HOX or PBX proteins to form dimers , or to a DNA-bound dimer of PBX and HOX proteins and thought to have a role in stabilization of the homeoprotein-DNA complex. Isoform 3 is required for the activity of a PDX1:PBX1b:MEIS2b complex in pancreatic acinar cells involved in the transcriptional activation of the ELA1 enhancer; the complex binds to the enhancer B element and cooperates with the transcription factor 1 complex (PTF1) bound to the enhancer A element; MEIS2 is not involved in complex DNA-binding. Probably in complex with PBX1 , is involved in transcriptional regulation by KLF4. Isoform 3 and isoform 4 can bind to a EPHA8 promoter sequence containing the DNA motif 5'-CGGTCA-3'; in cooperation with a PBX protein (such as PBX2) is proposed to be involved in the transcriptional



activation of EPHA8 in the developing midbrain. May be involved in regulation of myeloid differentiation. Can bind to the DNA sequence 5'-TGACAG-3' in the activator ACT sequence of the D (1A) dopamine receptor (DRD1) promoter and activate DRD1 transcription; isoform 5 cannot activate DRD1 transcription.

Background

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images