



SRC-1 (Phospho Thr1179) rabbit pAb

Catalog No	YP-Ab-10426
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB; ELISA
Gene Name	NCOA1 BHLHE74 SRC1
Immunogen	Synthesized peptide derived from human SRC-1 (Phospho Thr1179)
Specificity	This antibody detects endogenous levels of Human,Mouse SRC-1 (Phospho Thr1179)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	bHLHe74;Class E basic helix-loop-helix protein 74;F SRC 1;Hin 2 protein;Hin2 protein;MGC129719;MGC12972;mNRC 1;NCoA 1;NCoA-1;Ncoa1; NCOA1_HUMAN;Nuclear receptor coactivator 1;Nuclear receptor coactivator protein 1;NY REN 52 antigen;Protein Hin 2;Protein Hin-2;Protein Hin2;Renal carcinoma antigen NY REN 52;Renal carcinoma antigen NY-REN-52;RIP 16; RIP16;SRC 1;SRC-1;Steroid receptor coactivator 1;NCOA1 BHLHE74 SRC1; SRC-1 (Phospho Thr1179);Nuclear receptor coactivator 1 (NCoA-1;EC 2.3.1.48; RIP160;SRC-1)
Observed Band	158kD
Calculated Molecular Weight	2 kDa; 157kD
Cell Pathway	Nucleus .
Tissue Specificity	Widely expressed.
Function	catalytic activity:Acetyl-CoA + histone = CoA + acetylhistone.,disease:A chromosomal aberration involving NCOA1 is a cause of rhabdomyosarcoma. Translocation t(2;2)(q35;p23) with PAX3 generates the NCOA1-PAX3 oncogene consisting of the N-terminus part of PAX3 and the C-terminus part of NCOA1. The fusion protein acts as a transcriptional activator. Rhabdomyosarcoma is the



most common soft tissue carcinoma in childhood, representing 5-8% of all malignancies in children.,domain:Contains 7 Leu-Xaa-Xaa-Leu-Leu (LXXLL) motifs. LXXLL motifs 3, 4 and 5 are essential for the association with nuclear receptors. LXXLL motif 7, which is not present in isoform 2, increases the affinity for steroid receptors in vitro.,domain:The C-terminal (1107-1441) part mediates the histone acetyltransferase (HAT) activity.,function:Nuclear receptor coactivator that directly binds nuclear receptors and stimulates t

Background

The protein encoded by this gene acts as a transcriptional coactivator for steroid and nuclear hormone receptors. It is a member of the p160/steroid receptor coactivator (SRC) family and like other family members has histone acetyltransferase activity and contains a nuclear localization signal, as well as bHLH and PAS domains. The product of this gene binds nuclear receptors directly and stimulates the transcriptional activities in a hormone-dependent fashion. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images