



NRF2(Phospho-Ser40) rabbit pAb

Catalog No	YP-Ab-10437
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	NFE2L2 NRF2
Immunogen	Synthesized peptide derived from human NRF2(Phospho-Ser40)
Specificity	This antibody detects endogenous levels of NRF2(Phospho-Ser40) at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.86% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	erythroid derived 2;HEBP1;like 2;NF E2 related factor 2;NF-E2-related factor 2; NF2L2_HUMAN;NFE2 related factor 2;NFE2-related factor 2;Nfe2l2;Nrf 2;NRF2; Nuclear factor (erythroid derived 2) like 2;Nuclear factor;nuclear factor erythroid 2 like 2;Nuclear factor erythroid 2 related factor 2;Nuclear factor erythroid 2-related factor 2;Nuclear factor erythroid derived 2 like 2;NFE2L2 NRF2; NRF2(Phospho-Ser40);Nuclear factor erythroid 2-related factor 2 (NF-E2-related factor 2) (NFE2-related factor 2) (HEBP1) (Nuclear factor;like 2)
Observed Band	75-100kD
Calculated Molecular Weight	1~12kD; 68kD
Cell Pathway	Cytoplasm, cytosol . Nucleus . Cytosolic under unstressed conditions: ubiquitinated and degraded by the BCR(KEAP1) E3 ubiquitin ligase complex (PubMed:15601839, PubMed:21196497). Translocates into the nucleus upon induction by electrophilic agents that inactivate the BCR(KEAP1) E3 ubiquitin ligase complex (PubMed:21196497). .
Tissue Specificity	Widely expressed. Highest expression in adult muscle, kidney, lung, liver and in fetal muscle.



Function

domain:Acidic activation domain in the N-terminus, and DNA binding domain in the C-terminus.,function:Transcription activator that binds to antioxidant response (ARE) elements in the promoter regions of target genes. Important for the coordinated up-regulation of genes in response to oxidative stress. May be involved in the transcriptional activation of genes of the beta-globin cluster by mediating enhancer activity of hypersensitive site 2 of the beta-globin locus control region.,PTM:Phosphorylation of Ser-40 by PKC in response to oxidative stress dissociates NFE2L2 from its cytoplasmic inhibitor KEAP1, promoting its translocation into the nucleus.,similarity:Belongs to the bZIP family.,similarity:Belongs to the bZIP family. CNC subfamily.,similarity:Contains 1 bZIP domain.,subcellular location:Cytosolic under unstressed conditions, translocates into the nucleus upon induction by electr

Background

This gene encodes a transcription factor which is a member of a small family of basic leucine zipper (bZIP) proteins. The encoded transcription factor regulates genes which contain antioxidant response elements (ARE) in their promoters; many of these genes encode proteins involved in response to injury and inflammation which includes the production of free radicals. Multiple transcript variants encoding different isoforms have been characterized for this gene. [provided by RefSeq, Sep 2015],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images