



USF1 (Phospho-Thr153) rabbit pAb

Catalog No	YP-Ab-10517
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB
Gene Name	USF1 BHLHB11 USF
Immunogen	Synthesized peptide derived from human USF1 (Phospho-Thr153)
Specificity	This antibody detects endogenous levels of USF1 (Phospho-Thr153) at Human, Mouse, Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.173% sodium azide.
Source	Polyclonal, Rabbit, IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	bHLHb11; FCHL; FCHL1; HYPLIP1; MLTF; MLTF1; UEF; Upstream stimulatory factor 1; USF; USF1; USF1 BHLHB11 USF; USF1 (Phospho-Thr153); Upstream stimulatory factor 1 (Class B basic helix-loop-helix protein 11) (bHLHb11) (Major late transcription factor 1)
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Nucleus.
Tissue Specificity	Kidney,
Function	disease: Genetic variations in USF1 are associated with combined hyperlipidemia type 1 (HYPLIP1) [MIM:602491]; also known as familial combined hyperlipidemia type 1 (FCHL1). HYPLIP1 is characterized by elevated levels of serum total cholesterol, triglycerides or both, and is observed in about 20% of individuals with premature coronary heart disease., function: Transcription factor that binds to a symmetrical DNA sequence (E-boxes) (5'-CACGTG-3') that is found in a variety of viral and cellular promoters., similarity: Contains 1 basic helix-loop-helix (bHLH) domain., subunit: Efficient DNA binding requires dimerization with another bHLH protein. Binds DNA as a homodimer or a



heterodimer (USF1/USF2). Interacts with varicella-zoster virus IE62 protein.,

Background

This gene encodes a member of the basic helix-loop-helix leucine zipper family, and can function as a cellular transcription factor. The encoded protein can activate transcription through pyrimidine-rich initiator (Inr) elements and E-box motifs. This gene has been linked to familial combined hyperlipidemia (FCHL). Alternative splicing of this gene results in multiple transcript variants. A related pseudogene has been defined on chromosome 21. [provided by RefSeq, Feb 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of 1 HeLa cell, 2 LPS 100ng/mL 30min treated ,using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000