



NEC1 rabbit pAb

Catalog No	YP-Ab-11039
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB
Gene Name	PCSK1 NEC1
Immunogen	Synthesized peptide derived from human NEC1 AA range: 125-175
Specificity	This antibody detects endogenous levels of NEC1 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit, IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	BMIQ12; NEC 1; NEC1; Neuroendocrine convertase 1; PC1; PC3; PCSK1; Prohormone convertase 1; Proprotein; PCSK1 NEC1
Observed Band	
Calculated Molecular Weight	83kD
Cell Pathway	Cytoplasmic vesicle, secretory vesicle. Localized in the secretion granules.
Tissue Specificity	
Function	catalytic activity: Release of protein hormones, neuropeptides and renin from their precursors, generally by hydrolysis of -Lys-Arg- bonds., cofactor: Calcium., disease: Defects in PCSK1 are the cause of proprotein convertase 1 deficiency (PC1 deficiency) [MIM:600955]. PC1 deficiency is characterized by obesity, hypogonadism, hypoadrenalism, reactive hypoglycemia as well as marked small-intestinal absorptive dysfunction It is due to impaired processing of prohormones., function: Involved in the processing of hormone and other protein precursors at sites comprised of pairs of basic amino acid residues. Substrates include POMC, renin, enkephalin, dynorphin, somatostatin and insulin., polymorphism: Genetic variations in PCSK1 define the body mass index quantitative trait locus 12 (BMIQ12) [MIM:612362]. Variance in body mass index is a susceptibility factor for obesity., similarity: Belongs to the pe



Background

This gene encodes a member of the subtilisin-like proprotein convertase family, which includes proteases that process protein and peptide precursors trafficking through regulated or constitutive branches of the secretory pathway. The encoded protein undergoes an initial autocatalytic processing event in the ER to generate a heterodimer which exits the ER and sorts to subcellular compartments where a second autocatalytic even takes place and the catalytic activity is acquired. The protease is packaged into and activated in dense core secretory granules and expressed in the neuroendocrine system and brain. This gene encodes one of the seven basic amino acid-specific members which cleave their substrates at single or paired basic residues. It functions in the proteolytic activation of polypeptide hormones and neuropeptides precursors. Mutations in this gene have been associated with susceptibility to obesity and proprotein convertase 1/3 deficiency. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene [provided by RefSeq, Jan 2014],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from 293T cells, primary antibody was diluted at 1:1000, 4° over night