



MPPB rabbit pAb

Catalog No	YP-Ab-11094
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB; IHC
Gene Name	PMPCB MPPB
Immunogen	Synthesized peptide derived from human MPPB AA range: 92-142
Specificity	This antibody detects endogenous levels of MPPB at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit, IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000; IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Beta MPP; MPP11; MPPB; MPPP52; P 52; PMPCB; PMPCB MPPB
Observed Band	54kD
Calculated Molecular Weight	54kD
Cell Pathway	Mitochondrion matrix .
Tissue Specificity	
Function	catalytic activity: Release of N-terminal transit peptides from precursor proteins imported into the mitochondrion, typically with Arg in position P2., cofactor: Binds 1 zinc ion per subunit., function: Cleaves presequences (transit peptides) from mitochondrial protein precursors., similarity: Belongs to the peptidase M16 family., subunit: Heterodimer of alpha and beta subunits.,
Background	This gene is a member of the peptidase M16 family and encodes a protein with a zinc-binding motif. This protein is located in the mitochondrial matrix and catalyzes the cleavage of the leader peptides of precursor proteins newly imported into the mitochondria, though it only functions as part of a heterodimeric complex. [provided by RefSeq, Jul 2008],




matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from 293T cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 45min).