



FAIM3 rabbit pAb

Catalog No	YP-Ab-11234
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB
Gene Name	FAIM3 TOSO
Immunogen	Synthesized peptide derived from human FAIM3 AA range: 123-173
Specificity	This antibody detects endogenous levels of FAIM3 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit, IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	TOSO; FAIM3; FcμR; FAIM3 TOSO
Observed Band	
Calculated Molecular Weight	
Cell Pathway	[Isoform 1]: Membrane ; Single-pass membrane protein .; [Isoform 3]: Secreted . Detected in the serum. .
Tissue Specificity	Expressed in lymph nodes, peripheral blood leukocytes, lung, thymus and kidneys. Very weak expression detected in spleen, liver, heart, and salivary gland. Expressed in lymphoid cell lines such as Jurkat, CEM-T4, MOLT-4, HB11; 19 and Reh. No expression detected in nonhematopoietic cell lines including Hep-G2, HEK293 and HeLa. Detected at high levels in chronic lymphocytic leukemia cells.
Function	domain: The Ig-like domain is required for the anti-apoptotic ability. function: May play a role in the immune system processes. Protects cells from FAS-, TNF alpha- and FADD-induced apoptosis without increasing expression of the inhibitors of apoptosis BCL2 and BCLXL. Seems to activate an inhibitory pathway that prevents CASP8 activation following FAS stimulation, rather than blocking apoptotic signals downstream. May inhibit FAS-induced apoptosis by preventing CASP8 processing through CFLAR up-regulation. induction: By T-cell



activation.,miscellaneous:'Toso' is a Japanese liquor drunk on New Year's day to celebrate long life and eternal youth.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain.,tissue specificity:Expressed in lymph nodes, peripheral blood leukocytes, lung, thymus and kidneys. Very weak expression detected in spleen, liver, heart, and salivary gland. Expressed in I

Background

Fc receptors specifically bind to the Fc region of immunoglobulins (Igs) to mediate the unique functions of each Ig class. FAIM3 encodes an Fc receptor for IgM (see MIM 147020) (Kubagawa et al., 2009 [PubMed 19858324]; Shima et al., 2010 [PubMed 20042454]).[supplied by OMIM, Jul 2010],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4° over night