



NEUG rabbit pAb

Catalog No	YP-Ab-11831
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	IHC; IF
Gene Name	NRGN
Immunogen	Synthesized peptide derived from human NEUG AA range: 23-73
Specificity	This antibody detects endogenous levels of NEUG at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit, IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1:50-200. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Protein kinase C substrate RC3; Calmodulin binding protein; Hng; NEUG(55-78); NEUG_HUMAN; Neurogranin (protein kinase C substrate); Ng; NRGN; Protein kinase C substrate; RC3; NEUG
Observed Band	
Calculated Molecular Weight	
Cell Pathway	nucleus, trans-rk transport vesicle membrane, postsynaptic density, axon, mitochondrial membrane, neuronal cell body, dendritic spine head,
Tissue Specificity	In the cerebral cortex, found in the cell bodies of neurons in layers II-VI, and in apical and basal dendrites of pyramidal neurons. Is not found in the dendrites in patients with Alzheimer disease.
Function	function: Acts as a "third messenger" substrate of protein kinase C-mediated molecular cascades during synaptic development and remodeling. Binds to calmodulin in the absence of calcium., similarity: Belongs to the neurogranin family., similarity: Contains 1 collagen-like domain., similarity: Contains 1 IQ domain., tissue specificity: In the cerebral cortex, found in the cell bodies of neurons in layers II-VI, and in apical and basal dendrites of pyramidal neurons. Is not found in the dendrites in patients with Alzheimer disease.,



Background

Neurogranin (NRGN) is the human homolog of the neuron-specific rat RC3/neurogranin gene. This gene encodes a postsynaptic protein kinase substrate that binds calmodulin in the absence of calcium. The NRGN gene contains four exons and three introns. The exons 1 and 2 encode the protein and exons 3 and 4 contain untranslated sequences. It is suggested that the NRGN is a direct target for thyroid hormone in human brain, and that control of expression of this gene could underlay many of the consequences of hypothyroidism on mental states during development as well as in adult subjects. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Tris-EDTA, pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200 (4° overnight). 3, Secondary antibody was diluted at 1:200 (room temperature, 45min).