



ADCY5 rabbit pAb

Catalog No	YP-Ab-12075
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB
Gene Name	ADCY5
Immunogen	Synthesized peptide derived from human ADCY5 AA range: 471-521
Specificity	This antibody detects endogenous levels of ADCY5 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit, IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	AC5; Adenylate cyclase type V; Adenylate cyclase type 5; adenylate cyclase 5; ADCY5
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Cell membrane ; Multi-pass membrane protein. Cell projection, cilium .
Tissue Specificity	Detected in pancreas islets (at protein level). Detected in pancreas islets.
Function	catalytic activity: ATP = 3',5'-cyclic AMP + diphosphate., cofactor: Binds 2 magnesium ions per subunit., enzyme regulation: Inhibition by calcium in the submicromolar concentration range., function: This is a membrane-bound, calcium-inhibitable adenylyl cyclase., similarity: Belongs to the adenylyl cyclase class-4/guanylyl cyclase family., similarity: Contains 2 guanylate cyclase domains.,
Background	This gene encodes a member of the membrane-bound adenylyl cyclase enzymes. Adenylyl cyclases mediate G protein-coupled receptor signaling through the synthesis of the second messenger cAMP. Activity of the encoded protein is stimulated by the Gs alpha subunit of G protein-coupled receptors and is inhibited by protein kinase A, calcium and Gi alpha subunits. Single nucleotide



polymorphisms in this gene may be associated with low birth weight and type 2 diabetes. Alternatively spliced transcript variants that encode different isoforms have been observed for this gene. [provided by RefSeq, Dec 2010],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4° over night