



EP400 rabbit pAb

Catalog No	YP-Ab-12128
Isotype	IgG
Reactivity	Human; Mouse
Applications	IHC;IF
Gene Name	EP400 CAGH32 KIAA1498 KIAA1818 TNRC12
Immunogen	Synthesized peptide derived from human EP400 AA range: 2057-2107
Specificity	This antibody detects endogenous levels of EP400 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1:50-200. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CAGH32;E1A-binding protein p400;EC:3.6.4.-;EP 400;hDomino;EP400;EP400 CAGH32 KIAA1498 KIAA1818 TNRC12
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Nucleus .
Tissue Specificity	Ubiquitously expressed.
Function	caution:The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data., function:Component of the NuA4 histone acetyltransferase complex which is involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A. This modification may both alter nucleosome - DNA interactions and promote interaction of the modified histones with other proteins which positively regulate transcription. May be required for transcriptional activation of E2F1 and MYC target genes during cellular proliferation. The NuA4 complex ATPase and helicase activities seem to be, at least in part, contributed by the association of RUVBL1 and RUVBL2 with EP400. May regulate ZNF42 transcription activity.,sequence caution:Contaminating sequence. Potential poly-A sequence.,sequence

caution: Intron retention., similar

Background


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 45min).