



DAD1 rabbit pAb

Catalog No	YP-Ab-12234
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB
Gene Name	DAD1
Immunogen	Synthesized peptide derived from human DAD1 AA range: 38-88
Specificity	This antibody detects endogenous levels of DAD1 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit, IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Dolichyl-diphosphooligosaccharide--protein glycosyltransferase subunit DAD1; DAD-1; DAD1
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Endoplasmic reticulum membrane; Multi-pass membrane protein .
Tissue Specificity	
Function	catalytic activity:Dolichyl diphosphooligosaccharide + protein L-asparagine = dolichyl diphosphate + a glycoprotein with the oligosaccharide chain attached by N-glycosyl linkage to protein L-asparagine.,function:Component of the N-oligosaccharyl transferase enzyme which catalyzes the transfer of a high mannose oligosaccharide from a lipid-linked oligosaccharide donor to an asparagine residue within an Asn-X-Ser/Thr consensus motif in nascent polypeptide chains. N-glycosylation occurs cotranslationally and the complex associates with the Sec61 complex at the channel-forming translocon complex that mediates protein translocation across the endoplasmic reticulum (ER). Loss of the DAD1 protein triggers apoptosis.,similarity:Belongs to the DAD/OST2 family.,subunit:Component of the oligosaccharyltransferase (OST) complex. OST seems to exist in different forms which contain at least RPN1, RPN2,



Background

DAD1, the defender against apoptotic cell death, was initially identified as a negative regulator of programmed cell death in the temperature sensitive tsBN7 cell line. The DAD1 protein disappeared in temperature-sensitive cells following a shift to the nonpermissive temperature, suggesting that loss of the DAD1 protein triggered apoptosis. DAD1 is believed to be a tightly associated subunit of oligosaccharyltransferase both in the intact membrane and in the purified enzyme, thus reflecting the essential nature of N-linked glycosylation in eukaryotes. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from MDA-MB cells, primary antibody was diluted at 1:1000, 4° over night