



DREB rabbit pAb


Catalog No	YP-Ab-12300
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB
Gene Name	DBN1 D0S117E
Immunogen	Synthesized peptide derived from human DREB AA range: 484-534
Specificity	This antibody detects endogenous levels of DREB at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit, IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	DBN1; DBN 1; Developmentally-regulated brain protein; drebrin 1; DREB; DBN1 D0S117E
Observed Band	
Calculated Molecular Weight	
Cell Pathway	Cytoplasm . Cell projection, dendrite . Cytoplasm, cell cortex . Cell junction . Cell projection, growth cone . In the absence of antigen, evenly distributed throughout subcortical regions of the T-cell membrane and cytoplasm (PubMed:20215400). In the presence of antigen, distributes to the immunological synapse forming at the T-cell-APC contact area, where it localizes at the peripheral and distal supramolecular activation clusters (SMAC) (PubMed:20215400). Colocalized with RUFY3 and F-actin at the transitional domain of the axonal growth cone (By similarity). .
Tissue Specificity	Expressed in the brain, with expression in the molecular layer of the dentate gyrus, stratum pyramidale, and stratum radiatum of the hippocampus (at protein level) (PubMed:8838578). Also expressed in the terminal varicosities distributed along dendritic trees of pyramidal cells in CA4 and CA3 of the hippocampus (at protein level) (PubMed:8838578). Expressed in pyramidal cells in CA2, CA1 and the subiculum of the hippocampus (at protein level) (PubMed:8838578). Expressed in peripheral blood lymphocytes, including T-cells (at protein level)



(PubMed:20215400). Expressed in the brain (PubMed:8216329, Ref.2). Expressed in the heart, placenta, lung, skeletal muscle, kidney, pancreas, skin fibroblasts, gingival fibroblasts and bone-derived cells (Ref.2).

Function	function:Drebrins might play some role in cell migration, extension of neuronal processes and plasticity of dendrites, respectively.,similarity:Contains 1 ADF-H domain.,subunit:Binds F-actin.,tissue specificity:Brain neurons. Also found in the heart, placenta, skeletal muscle, kidney and pancreas.,
Background	The protein encoded by this gene is a cytoplasmic actin-binding protein thought to play a role in the process of neuronal growth. It is a member of the drebrin family of proteins that are developmentally regulated in the brain. A decrease in the amount of this protein in the brain has been implicated as a possible contributing factor in the pathogenesis of memory disturbance in Alzheimer's disease. At least two alternative splice variants encoding different protein isoforms have been described for this gene. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HeLa cells, primary antibody was diluted at 1:1000, 4° over night