



OPTN rabbit pAb

Catalog No	YP-Ab-12586
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	OPTN FIP2 GLC1E HIP7 HYPL NRP
Immunogen	Synthesized peptide derived from human OPTN
Specificity	This antibody detects endogenous levels of OPTN at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.78% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Optineurin;OPTN;ALS12;E3-14.7K-interacting protein;FAA4;OPTN FIP2 GLC1E HIP7 HYPL NRP;Optineurin (E3-14.7K-interacting protein) (FIP-2) (Huntingtin yeast partner L) (Huntingtin-interacting protein 7) (HIP-7) (Huntingtin-interacting protein L) (NEMO-related protein) (Optic neuropathy-inducing protein) (Transcription factor IIIA-interacting protein) (TFIIIA-IntP)
Observed Band	66-70 kDa
Calculated Molecular Weight	66 kDa
Cell Pathway	Cytoplasm, perinuclear region. Golgi apparatus . Golgi apparatus, trans-Golgi network. Cytoplasmic vesicle, autophagosome. Cytoplasmic vesicle. Recycling endosome. Found in the perinuclear region and associates with the Golgi apparatus (PubMed:27534431). Colocalizes with MYO6 and RAB8 at the Golgi complex and in vesicular structures close to the plasma membrane. Localizes to LC3-positive cytoplasmic vesicles upon induction of autophagy. .
Tissue Specificity	Present in aqueous humor of the eye (at protein level). Highly expressed in trabecular meshwork. Expressed nonpigmented ciliary epithelium, retina, brain, adrenal cortex, fetus, lymphocyte, fibroblast, skeletal muscle, heart, liver, brain and placenta.
Function	

Background

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images