



GAP-43 Monoclonal Antibody(Mix)

Catalog No	YP-Ab-12628
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;IHC;IF
Gene Name	GAP43
Immunogen	Recombinant Protein of GAP-43
Specificity	The antibody detects endogenous GAP-43 protein.
Formulation	PBS, pH 7.4, containing 0.5%BSA, 0.02% sodium azide as Preservative and 50% Glycerol.
Source	Monoclonal, Mouse
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using epitope-specific immunogen.
Dilution	WB: 1:1000-2000 IHC:1:200-500. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Axonal membrane protein GAP 43;Axonal membrane protein GAP-43;B 50; growth associated protein 43;GAP-43;GAP43;Neuromodulin;Growth-associated protein 43;Neural phosphoprotein B-50;pp46
Observed Band	38,43kD
Calculated Molecular Weight	38,43kD
Cell Pathway	Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cell projection, growth cone membrane ; Peripheral membrane protein ; Cytoplasmic side . Cell junction, synapse . Cell projection, filopodium membrane ; Peripheral membrane protein . Perikaryon . Cell projection, dendrite . Cell projection, axon . Cytoplasm . Cytoplasmic surface of growth cone and synaptic plasma membranes. .
Tissue Specificity	Alzheimer cortex,Brain,Subthalamic nucleus,
Function	function:This protein is associated with nerve growth. It is a major component of the motile "growth cones" that form the tips of elongating axons.,online information:Gap-43 entry,PTM:Phosphorylation of this protein by a protein kinase C is specifically correlated with certain forms of synaptic plasticity., similarity:Belongs to the neuromodulin family.,similarity:Contains 1 IQ domain.,



subcellular location: Cytoplasmic surface of growth cone and synaptic plasma membranes., subunit: Binds calmodulin with a greater affinity in the absence of Ca^{2+} than in its presence.,

Background

The protein encoded by this gene has been termed a 'growth' or 'plasticity' protein because it is expressed at high levels in neuronal growth cones during development and axonal regeneration. This protein is considered a crucial component of an effective regenerative response in the nervous system. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of 1) Hela, 2) 293T, 3) Mouse Brain Tissue, 4) Rat Brain Tissue using GAP-43 Monoclonal Antibody.



Immunohistochemical analysis of paraffin-embedded Rat Brain Tissue using GAP-43 Monoclonal Antibody.