



Synuclein-α (phospho Tyr136) Polyclonal Antibody


Catalog No	YP-Ab-12644
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	SNCA
Immunogen	The antiserum was produced against synthesized peptide derived from human Synuclein-alpha around the phosphorylation site of Tyr136. AA range:91-140
Specificity	Phospho-Synuclein-α (Y136) Polyclonal Antibody detects endogenous levels of Synuclein-α protein only when phosphorylated at Y136.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Alpha synuclein;Alpha-synuclein;isoform NACP14;alphaSYN;MGC15443;MGC11988;MGC12756;MGC64356;NACP;Non A beta component of AD amyloid;Non A4 component of amyloid;Non A4 component of amyloid precursor;Non-A beta component of AD amyloid;Non-A-beta component of alzheimers disease amyloid;precursor of;Non-A4 component of amyloid precursor;Non-A4 component of amyloid;OTTHUMP218549;OTTHUMP218551;OTTHUMP218552;OTTHUMP218553;OTTHUMP218554;PARK 1;PARK 4;PARK1;PARK4;Parkinson disease (autosomal dominant;Lewy body) 4;Parkinson disease familial 1;SNCA;Snca synuclein;alpha (non A4 component of amyloid precursor);SYN;Synuclein alpha;Synuclein alpha 14;Synuclein;SYUA_HUMAN;Synuclein-α
Observed Band	15kD
Calculated Molecular Weight	15kD
Cell Pathway	Cytoplasm . Membrane . Nucleus . Cell junction, synapse . Secreted . Cell projection, axon . Membrane-bound in dopaminergic neurons (PubMed:15282274). Expressed and colocalized with SEPTIN4 in dopaminergic axon terminals, especially at the varicosities (By similarity). .
Tissue Specificity	Highly expressed in presynaptic terminals in the central nervous system. Expressed principally in brain.




Function	alternative products: Additional isoforms seem to exist, disease: Brain iron accumulation type 1 (NBIA1, also called Hallervorden-Spatz syndrome), a rare neuroaxonal dystrophy, is histologically characterized by axonal spheroids, iron deposition, Lewy body (LB)-like intraneuronal inclusions, glial inclusions and neurofibrillary tangles. SNCA is found in LB-like inclusions, glial inclusions and spheroids., disease: Defects in SNCA are a cause of autosomal dominant Parkinson disease 1 (PARK1) [MIM:168601, 168600]. Parkinson disease (PD) is a complex, multifactorial disorder that typically manifests after the age of 50 years, although early-onset cases (before 50 years) are known. PD generally arises as a sporadic condition but is occasionally inherited as a simple mendelian trait. Although sporadic and familial PD are very similar, inherited forms of the disease usually begin at earlier ages an
Background	Alpha-synuclein is a member of the synuclein family, which also includes beta- and gamma-synuclein. Synucleins are abundantly expressed in the brain and alpha- and beta-synuclein inhibit phospholipase D2 selectively. SNCA may serve to integrate presynaptic signaling and membrane trafficking. Defects in SNCA have been implicated in the pathogenesis of Parkinson disease. SNCA peptides are a major component of amyloid plaques in the brains of patients with Alzheimer's disease. Alternatively spliced transcripts encoding different isoforms have been identified for this gene. [provided by RefSeq, Feb 2016],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Enzyme-Linked Immunosorbent Assay
(Phospho-ELISA) for Immunogen Phosphopeptide
(Phospho-left) and Non-Phosphopeptide
(Phospho-right), using Synuclein-alpha
(Phospho-Tyr136) Antibody



Western blot analysis of lysates from mouse brain,
using Synuclein-alpha (Phospho-Tyr136) Antibody.
The lane on the right is blocked with the phospho
peptide.