




Trk A (Phospho-Y496) Antibody

Catalog No	YP-Ab-12666
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB; IF; ELISA
Gene Name	NTRK1 MTC TRK TRKA
Immunogen	Synthesized pospho peptide derived from human Trk A (Phospho-Y496)
Specificity	This antibody detects endogenous pospho levels of human Trk A (Phospho-Y496)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from mouse ascites by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000; IF/ICC 1:100-500; ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	gp14trk;High affinity nerve growth factor receptor;High affinity nerve growth factor receptor precursor;MTC;Neurotrophic tyrosine kinase receptor type 1; NTRK1;NTRK1_HUMAN;Oncogene TRK;p14-TrkA;p14 TrkA;Slow nerve growth; Trk A;TRK;Trk-A;TRK1;TRK1-transforming tyrosine kinase protein;Tropomyosin-related kinase A;Tyrosine kinase receptor A;Tyrosine kinase receptor;NTRK1 MTC TRK TRKA;Trk A (Phospho-Y496);High affinity nerve growth factor receptor (EC 2.7.10.1;gp140trk;p140-TrkA)
Observed Band	180kD
Calculated Molecular Weight	180kD
Cell Pathway	Cell membrane ; Single-pass type I membrane protein . Early endosome membrane ; Single-pass type I membrane protein . Late endosome membrane ; Single-pass type I membrane protein . Recycling endosome membrane ; Single-pass type I membrane protein . Rapidly internalized after NGF binding (PubMed:1281417). Internalized to endosomes upon binding of NGF or NTF3 and further transported to the cell body via a retrograde axonal transport. Localized at cell membrane and early endosomes before nerve growth factor (NGF) stimulation. Recruited to late endosomes after NGF stimulation. Colocalized with RAPGEF2 at late endosomes. .



Tissue Specificity	Isoform TrkA-I is found in most non-neuronal tissues. Isoform TrkA-II is primarily expressed in neuronal cells. TrkA-III is specifically expressed by pluripotent neural stem and neural crest progenitors.
Function	alternative products: Both isoforms have similar biological properties, catalytic activity: $\text{ATP} + \text{a [protein]-L-tyrosine} = \text{ADP} + \text{a [protein]-L-tyrosine phosphate.}$, caution: The sequence shown here is derived from an Ensembl automatic analysis pipeline and should be considered as preliminary data., disease: Chromosomal aberrations involving NTRK1 are a cause of thyroid papillary carcinoma (PACT) [MIM:188550]. Intrachromosomal rearrangement that links the protein kinase domain of NTRK1 to the 5'-end of the TPR gene forms the fusion protein TRK-T1. TRK-T1 is a 55 kDa protein reacting with antibodies against the C-terminus of the NTRK1 protein., disease: Chromosomal aberrations involving NTRK1 are a cause of thyroid papillary carcinoma (PACT) [MIM:188550]. Translocation t(1;3)(q21;q11) with TFG generates the TRKT3 (TRK-T3) transcript by fusing TFG to the 3'-end of NTRK1; a rearrangement with TPM3 gen
Background	This gene encodes a member of the neurotrophic tyrosine kinase receptor (NTRK) family. This kinase is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. The presence of this kinase leads to cell differentiation and may play a role in specifying sensory neuron subtypes. Mutations in this gene have been associated with congenital insensitivity to pain, anhidrosis, self-mutilating behavior, mental retardation and cancer. Alternate transcriptional splice variants of this gene have been found, but only three have been characterized to date. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of various lysates, primary antibody was diluted at 1:1000, 4° over night, secondary antibody(cat: RS23920)was diluted at 1:10000, 37° 1hour.